

## DATA SHEET

### NAME OF MEDICINE

#### RISPERDAL®

Risperidone

0.5 mg, 1 mg, 2 mg, 3 mg, 4 mg tablets

0.5 mg, 1 mg, 2 mg, 3 mg, 4 mg Quicklet® orally-disintegrating tablets

1 mg/mL oral solution

### PRESENTATION

RISPERDAL is available as oral, film-coated tablets of the following strengths:

- |                     |   |
|---------------------|---|
| 0.5 mg risperidone: | brownish-red, film-coated biconvex, half-scored oblong tablets<br>(marked on the grooved side with "Ris 0.5" and "JANSSEN" on the other side) |
| 1 mg risperidone:   | white, film-coated, half-scored, oblong tablets<br>(marked on the grooved side with "Ris 1")  |
| 2 mg risperidone:   | orange, film-coated, half-scored, oblong tablets<br>(marked on the grooved side with "Ris 2")   |
| 3 mg risperidone:   | yellow, film-coated, half-scored, oblong tablets<br>(marked on the grooved side with "Ris 3")   |
| 4 mg risperidone:   | green, film-coated, half-scored, oblong tablets<br>(marked on the grooved side with "Ris 4")  |

RISPERDAL is also available as a 1 mg/mL oral solution

RISPERDAL Quicklet is available as orally-disintegrating tablets of the following strengths:

- |                     |  |
|---------------------|--|
| 0.5 mg risperidone: | round, light coral, biconvex, etched on one side with "R0.5" |
| 1 mg risperidone:   | square, light coral, biconvex, etched on one side with "R1"  |
| 2 mg risperidone:   | Square, coral, biconvex, etched on one side with "R2"        |
| 3 mg risperidone:   | Round, coral, biconvex, etched on one side with "R3"         |
| 4 mg risperidone:   | Round, coral, biconvex, etched on one side with "R4"         |

### USES

#### *Actions*

Risperidone is a compound, which belongs to a new class of antipsychotic agents, the benzisoxazole derivatives.

Risperidone is a selective monoaminergic antagonist having a high affinity for serotonergic 5-HT<sub>2</sub> and dopaminergic D<sub>2</sub> receptors. Risperidone binds also to alpha<sub>1</sub>-adrenergic receptors and, with lower affinity, to H<sub>1</sub>-histamine and alpha<sub>2</sub>-adrenergic receptors. Risperidone has no affinity for cholinergic receptors. The antipsychotic activity of risperidone is considered to be attributable to both risperidone and its active metabolite 9-hydroxy risperidone. Risperidone, as a potent D<sub>2</sub> antagonist, improves the positive symptoms of schizophrenia but causes less depression of motor activity and induction of catalepsy than classical neuroleptics. Balanced central serotonin and dopamine antagonism may reduce extrapyramidal side effect liability and extend the therapeutic activity to the negative and affective symptoms of schizophrenia.

### **Pharmacokinetics**

Risperidone is completely absorbed after oral administration, reaching peak plasma concentrations within 1 to 2 hours. The absorption is not affected by food and thus RISPERDAL may be given with or without meals.

Risperidone is partly metabolised by CYP2D6 to 9-hydroxyrisperidone, which has similar pharmacological activity to risperidone. Another metabolic pathway is N-dealkylation.

After oral administration to psychotic patients, risperidone is eliminated with a half-life of about 3 hours. The elimination half-life of 9-hydroxy-risperidone and of risperidone plus 9-hydroxy risperidone is 24 hours.

Steady-state of risperidone is reached within 1 day in most patients. Steady-state of 9-hydroxy-risperidone is reached within 4-5 days of dosing. Risperidone plasma concentrations are dose-proportional within the therapeutic dose range.

Risperidone is rapidly distributed. The volume of distribution is 1-2 L/kg. In plasma, risperidone is bound to albumin and alpha<sub>1</sub>-acid glycoprotein. The plasma protein binding of risperidone is 88%, while that of 9-hydroxy-risperidone is 77%.

One week after administration, 70% of the dose is excreted in the urine and 14% in the faeces. In urine, risperidone plus 9-hydroxy-risperidone represents 35-45% of the dose.

A single-dose study showed higher active plasma concentrations and a reduced clearance of risperidone plus 9-hydroxy risperidone by 30% in the elderly and 60% in patients with renal insufficiency. Risperidone plasma concentrations were normal in patients with liver insufficiency, but the unbound risperidone in plasma was increased by about 35%.

Risperidone Quicklet tablets are bioequivalent to conventional risperidone tablets.

## **INDICATIONS**

RISPERDAL is indicated for the treatment of schizophrenia and other psychotic disorders. These include first episode psychoses, acute schizophrenic exacerbations, chronic schizophrenia and other psychotic conditions, in which positive symptoms (such as hallucinations, delusions, thought disturbances, hostility, suspiciousness), and/or negative symptoms (such as blunted affect, emotional and social withdrawal, poverty of speech) are prominent.

RISPERDAL is also indicated for the treatment and long term control of mania in bipolar disorder. These episodes are characterised by symptoms such as elevated, expansive or irritable mood, inflated self esteem, decreased need for sleep, pressured speech, racing thoughts, distractibility, or poor judgement, including disruptive or aggressive behaviours.

RISPERDAL also alleviates affective symptoms (such as depression, guilt-feelings, anxiety) associated with schizophrenia. In addition, RISPERDAL also appears effective in maintaining the clinical improvement during continuation therapy in patients who have shown an initial response to treatment with this agent.

RISPERDAL is indicated for the treatment (up to 12 weeks) of agitation, aggression or psychotic symptoms in patients with moderate to severe dementia of the Alzheimer type.

RISPERDAL is also indicated for the treatment of conduct and other disruptive behaviour disorders in children (over 5 years), adolescents and adults with subaverage intellectual functioning or mental retardation, or average IQ, in whom destructive behaviours (e.g. aggression, impulsivity and self-injurious behaviours) are prominent. RISPERDAL is also effective in maintaining the clinical improvement during continuation therapy in children and adolescents who have shown an initial treatment response. Pharmacological treatment should be an integral part of a more comprehensive treatment program, including psychosocial and educational intervention. Treatment with RISPERDAL for patients with disruptive behaviour disorders should be initiated only in consultation with a specialist, including child and adolescent

psychiatrists, paediatric neurologists, developmental paediatricians, or other physicians conversant in the diagnosis and treatment of conduct and other disruptive behaviour disorders. RISPERSDAL is indicated for the treatment of autism in children and adolescents.

## **DOSAGE AND ADMINISTRATION**

RISPERSDAL may be given as tablets or oral solution.

### ***Schizophrenia***

#### Switching from other antipsychotics

When medically appropriate, gradual discontinuation of the previous treatment while RISPERSDAL therapy is initiated is recommended. Also if medically appropriate, when switching patients from depot antipsychotics, initiate RISPERSDAL therapy in place of the next scheduled injection. The need for continuing existing anti-Parkinson medications should be re-evaluated periodically.

#### Adults

RISPERSDAL may be given once daily or twice daily. Patients should start with 2 mg/day RISPERSDAL. The dose may be increased on the second day to 4 mg. From then on the dosage can be maintained unchanged, or further individualised, if needed. Most patients will benefit from daily doses between 4 and 6 mg. In some patients a slower titration phase and a lower starting and maintenance dose may be appropriate.

Doses above 10 mg/day have not been shown to be superior in efficacy to lower doses and may cause extrapyramidal symptoms. Since the safety of doses above 16 mg/day has not been evaluated, doses above this level should not be used.

A benzodiazepine may be added to RISPERSDAL when additional sedation is required.

#### Elderly

A starting dose of 0.5 mg twice daily is recommended. This dosage can be individually adjusted with 0.5 mg twice daily increments to 1 - 2 mg twice daily.

RISPERSDAL is well tolerated in the elderly.

#### Children

Experience is lacking in children aged less than 15 years.

### **Bipolar Mania**

RISPERSDAL should be administered on a once daily schedule, starting with 2mg. Dosage adjustments, if indicated, should occur at intervals of not less than 24 hours and in dosage increments of 1mg per day. A dosing range between 2 and 6mg per day is recommended. As with all symptomatic treatments, the continued use of RISPERSDAL must be evaluated and justified on an ongoing basis.

### **Behavioural Disturbances in Patients with Dementia**

A starting dose of 0.25 mg twice daily is recommended. This dosage can be individually adjusted by increments of 0.25 mg twice daily, not more frequently than every other day, if needed. The optimum dose is 0.5 mg twice daily for most patients. Some patients, however, may benefit from doses up to 1 mg twice daily.

Once patients have reached their target dose, a once daily dosing regimen can be considered. As with all symptomatic treatments, the continued use of RISPERSDAL must be evaluated and justified on an on-going basis.

## Conduct and other disruptive behaviour disorders

For Subjects  $\geq 50$ kg

A starting dose of 0.5mg once daily is recommended. This dosage can be individually adjusted by increments of 0.5mg once daily not more frequently than every other day, if needed. The optimum dose is 1mg once daily for most patients. Some patients, however, may benefit from 0.5mg once daily while others may require 1.5mg once daily.

For Subjects  $< 50$ kg

A starting dose of 0.25mg once daily is recommended, which can be individually adjusted by increments of 0.25mg once daily not more frequently than every other day, if needed. The optimum dose is 0.5mg once daily for most patients, although some patients may benefit from 0.25mg once daily while others may require 0.75mg once daily.

As with all symptomatic treatments, the continued use of RISPERDAL must be evaluated and justified on an on-going basis.

## Autism

RISPERDAL can be administered once or twice daily. Patients experiencing somnolence may benefit from a switch in dosing from once daily to either once daily at bedtime, or twice daily.

RISPERDAL should be administered based on body weight. Dosing should begin at 0.25 mg or 0.5 mg/day based upon weight (see table below for relative weight categories). On Day 4 of treatment, the dose may be increased up to 0.5 mg or 1.0mg/day. This dose should be maintained and response assessed at approximately day 14. Only in patients not achieving sufficient clinical response should additional dose increases be considered. Dose increases may proceed at  $\geq 2$ -week intervals in increments of 0.25 mg for patients  $< 20$  kg or 0.5 mg for patients  $\geq 20$  kg. Based upon current studies, the maximum dose studied did not exceed a total daily dose of 1.5 mg in patients  $< 20$  kg, 2.5 mg in patients  $\geq 20$  kg and 3.5 mg in patients  $> 45$  kg. Doses below 0.25 mg/day were not effective in clinical studies.

The table of the maximum daily doses provides a reference for titration and dosing by weight based upon current studies, and may serve as a guide according to clinical need:

<b>Doses of RISPERDAL in Paediatric Patients with Autistic Disorder</b>				
Weight Categories	Days 1 – 3	Days 4 – 14+	Increments if dose increases are needed	Dose Range
<b>Dose by Weight in mg/day</b>				
$< 20$ kg	0.25 mg	0.5 mg	+0.25 mg at $\geq 2$ week intervals	0.5 mg-1.5 mg
$\geq 20$ kg	0.5 mg	1.0 mg	+0.5 mg at $\geq 2$ week intervals	1.0 mg-2.5 mg*
<b>Dose Range in mg/kg/day</b>				
			Increments if dose increases are needed	Dose Range
All	0.01 mg/kg/d	0.02 mg/kg/d	+0.01 mg/kg/day at $\geq 2$ week intervals	0.02 mg/kg/d-0.06 mg/kg/d

\* Subjects weighing  $> 45$  kg may require higher doses: maximum dose studied was 3.5 mg/day

Once sufficient response has been achieved and maintained consideration may be given to gradually lowering the dose to achieve optimum balance of effectiveness and tolerance.

Clinical experience was limited in autistic adolescents and in autistic children with an IQ>84 as not many of these patients were included in the trials.

As with all symptomatic treatments, the continued use of RISPERDAL in children and adolescents with autism must be evaluated and justified on an ongoing basis.

### **Renal and Hepatic Impairment**

Patients with renal impairment have less ability to eliminate risperidone plus 9-hydroxy risperidone than normal adults. Patients with impaired hepatic function have increases in plasma concentration of unbound risperidone.

Irrespective of the indication, starting and consecutive dosing should be halved, and dose titration should be slower for patients with renal or hepatic impairment.

RISPERDAL should be used with caution in these groups of patients.

## **CONTRAINDICATIONS**

RISPERDAL is contraindicated in patients with a known hypersensitivity to the product.

## **WARNINGS AND PRECAUTIONS**

### ***Warnings***

#### ***Elderly Patients with Dementia***

##### **Overall Mortality**

Elderly patients with dementia treated with atypical antipsychotic drugs have an increased mortality compared to placebo in a meta-analysis of 17 controlled trials of atypical antipsychotic drugs, including RISPERDAL. In placebo-controlled trials with RISPERDAL in this population, the incidence of mortality was 4.0% (40/1009) for RISPERDAL treated patients compared to 3.1% (22/712) for placebo-treated patients. The mean age (range) of patients who died was 86 years (range 67-100).

##### **Concomitant use with Frusemide**

In the RISPERDAL placebo-controlled trials in elderly patients with dementia, a higher incidence of mortality was observed in patients treated with frusemide plus risperidone (7.3% [15/206]; mean age 89 years, range 75-97) when compared to patients treated with risperidone alone (3.1% [25/803]; mean age 84 years, range 70-96) or frusemide alone (4.1% [5/121]; mean age 80 years, range 67-90). The increase in mortality in patients treated with frusemide plus risperidone was observed in two of the four clinical trials.

No pathophysiological mechanism has been clearly identified to explain this finding, and no consistent pattern for cause of death was observed. Nevertheless, caution should be exercised and the risks and benefits of this combination should be considered prior to the decision to use. There was no increased mortality among patients taking other diuretics concomitantly with risperidone. Irrespective of treatment, dehydration was an overall risk factor for mortality and should therefore be carefully avoided in elderly patients with dementia.

## Cerebrovascular Adverse Events

In placebo-controlled trials in elderly patients with dementia, there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks in patients (mean age 85 years, range 73-97) treated with RISPERDAL compared with patients treated with placebo. The pooled data from six placebo-controlled studies in mainly elderly patients (>65 years of age) with dementia showed that cerebrovascular adverse events (serious and non-serious combined) occurred in 3.3% (33/989) of patients treated with risperidone and 1.2% (8/693) of patients treated with placebo. The Odds Ratio (95% exact confidence interval) was 2.96 (1.33, 7.45).

Risperidone may interfere with activities requiring mental alertness. Therefore, patients should be advised not to drive or operate machinery until their individual susceptibility is known.

## **Precautions**

### *Alpha blocking activity*

Due to the alpha-blocking activity of risperidone, orthostatic hypotension can occur, especially during the initial dose-titration period. Clinically significant hypotension has been observed postmarketing with concomitant use of risperidone and antihypertensive treatment. RISPERDAL should be used with caution in patients with known cardiovascular disease (eg. heart failure, myocardial infarction, conduction abnormalities, dehydration, hypovolaemia, or cerebrovascular disease), and the dosage should be gradually titrated as recommended (see **DOSAGE AND ADMINISTRATION**). A dose reduction should be considered if hypotension occurs. Special care should be taken in patients taking medications to lower blood pressure.

### *Tardive Dyskinesia*

Medicines with dopamine receptor antagonistic properties have been associated with the induction of tardive dyskinesia characterised by rhythmical involuntary movements, predominantly of the tongue and/or face. It has been reported that the occurrence of extrapyramidal symptoms is a risk factor for the development of tardive dyskinesia. Because RISPERDAL has a lower potential to induce extrapyramidal symptoms than classic neuroleptics, it should have a reduced risk of inducing tardive dyskinesia. If signs and symptoms of tardive dyskinesia appear, the discontinuation of all antipsychotic medicines should be considered.

### *Neuroleptic Malignant Syndrome*

The Neuroleptic Malignant Syndrome, characterised by hyperthermia, muscle rigidity, autonomic instability, altered consciousness and elevated creatine phosphokinase (CPK) levels has been reported to occur with classical neuroleptics. In this event, all antipsychotic medicines, including RISPERDAL, should be discontinued.

### *Parkinson's Disease and Dementia with Lewy Bodies*

Physicians should weigh the risks versus benefits when prescribing antipsychotics including RISPERDAL to patients with Parkinson's disease or Dementia with Lewy Bodies (DLB) since both groups may be at increased risk of Neuroleptic Malignant Syndrome as well as having an increased sensitivity to antipsychotic medications. Manifestation of this increased sensitivity can include confusion, obtundation, postural instability with frequent falls, in addition to extrapyramidal symptoms.

### *Hyperglycaemia and Diabetes Mellitus*

Hyperglycaemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including RISPERDAL. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycaemia related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycaemia-related adverse events in patients treated with atypical antipsychotics. Precise risk estimates for hyperglycaemia related adverse events in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g. obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycaemia including polydipsia, polyuria, polyphagia and weakness. Patients who develop symptoms of hyperglycaemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycaemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of antidiabetic treatment despite discontinuation of the suspect drug.

### *Leukopenia, Neutropenia, and Agranulocytosis*

Events of leukopenia, neutropenia, and agranulocytosis have been reported with antipsychotic agents, including RISPERDAL. Agranulocytosis has been reported very rarely (< 1/10,000 patients) during postmarketing surveillance.

Patients with a history of a clinically significant low white blood cell count (WBC) or a drug-induced leukopenia/neutropenia should be monitored during the first few months of therapy and discontinuation of RISPERDAL should be considered at the first sign of a clinically significant decline in WBC in the absence of other causative factors.

Patients with clinically significant neutropenia should be carefully monitored for fever or other symptoms or signs of infection and treated promptly if such symptoms or signs occur. Patients with severe neutropenia (absolute neutrophil count <  $1 \times 10^9/L$ ) should discontinue RISPERDAL and have their WBC followed until recovery.

### *Venous Thromboembolism*

Cases of venous thromboembolism (VTE) have been reported with antipsychotic drugs. Since patients treated with antipsychotics often present with acquired risk factors for VTE, all possible risk factors for VTE should be identified before and during treatment with RISPERDAL and preventive measures undertaken.

### *Priapism*

Drugs with alpha-adrenergic blocking effects have been reported to induce priapism. Priapism has been reported with RISPERDAL during postmarketing surveillance.

### *Body Temperature Regulation*

Disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Appropriate care is advised when prescribing RISPERDAL to patients who will be experiencing conditions which may contribute to an elevation in core body temperature, e.g. exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity, or being subject to dehydration.

### *Antiemetic Effect*

An antiemetic effect was observed in preclinical studies with risperidone. This effect, if it occurs in humans, may mask the signs and symptoms of overdose with certain drugs or of conditions such as intestinal obstruction, Reye's syndrome, and brain tumour.

### *Seizures*

As with other antipsychotic drugs, RISPERDAL should be used cautiously in patients with a history of seizures or other conditions that potentially lower seizure threshold.

### *Intraoperative Floppy Iris Syndrome*

Intraoperative Floppy Iris Syndrome (IFIS) has been observed during cataract surgery in patients treated with medicines with alpha<sub>1</sub>-adrenergic antagonist effect, including RISPERDAL (see **ADVERSE EFFECTS**).

IFIS may increase the risk of eye complications during and after the operation. Current or past use of medicines with alpha<sub>1</sub>-adrenergic antagonist effect should be made known to the ophthalmic surgeon in advance of surgery. The potential benefit of stopping alpha<sub>1</sub> blocking therapy prior to cataract surgery has not been established and must be weighed against the risk of stopping the antipsychotic therapy.

### *Weight Gain*

Significant weight gain has been reported. Monitoring weight gain is advisable when RISPERDAL is being used.

### *QT Interval*

As with other antipsychotics, caution should be exercised when RISPERDAL is prescribed in patients with a history of cardiac arrhythmias, in patients with congenital long QT syndrome, and in concomitant use with drugs known to prolong the QT interval.

### *Dysphagia*

Oesophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's dementia. RISPERDAL and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

### *Suicide*

The possibility of a suicide attempt is inherent in schizophrenia and bipolar disorder, and close supervision of high-risk patients should accompany therapy. Prescriptions for RISPERDAL should be written for the smallest quantity consistent with good patient management, in order to reduce the risk of overdose.

### *Akathisia*

The presentation of akathisia may be variable and comprises subjective complaints of restlessness and an overwhelming urge to move and either distress or motor phenomena such as pacing, swinging of the legs while seated, rocking from foot to foot, or both. Particular attention should be paid to the monitoring for such symptoms and signs as, left untreated, akathisia is associated with poor compliance and an increased risk of relapse.

### Use in the elderly

It is recommended to halve both the starting dose and the subsequent dose increments in geriatric patients.

### Use in renal impairment

It is recommended to halve both the starting dose and the subsequent dose increments in patients with renal insufficiency.

### Use in hepatic impairment

It is recommended to halve both the starting dose and the subsequent dose increments in patients with hepatic insufficiency.

### Pregnancy and Lactation

The safety of RISPERDAL for use during human pregnancy has not been established. Although, in experimental animals, risperidone did not show direct reproductive toxicity, some indirect, prolactin- and CNS-mediated effects were observed. No teratogenic effect of risperidone was noted in any study.

*Non-teratogenic class effect.* Neonates exposed to antipsychotic drugs (including RISPERDAL) during the third trimester of pregnancy are at risk of experiencing extrapyramidal neurological disturbances and/or withdrawal symptoms following delivery. There have been post-market reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeling disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited; in other cases neonates have required additional medical treatment or monitoring.

RISPERDAL should be used during pregnancy only if the anticipated benefit outweighs the risk and the administered dose and duration of treatment should be as low and as short as possible.

In animal studies, risperidone and 9-hydroxyrisperidone are excreted in the milk. It has been demonstrated that this excretion also occurs in human breast milk. Therefore, women receiving RISPERDAL should not breast feed.

## Paediatric Use

RISPERDAL had no adverse effects on cognitive function in paediatric patients. In combined, long-term, open-label trials, mean changes in cognitive function tests were small and did not increase or decrease over time.

A mean increase of 7.5 kg after 12 months of RISPERDAL treatment was observed, somewhat higher than the expected weight gain (approximately 3 to 3.5 kg per year) for children predominantly between 5 and 12 years of age.

RISPERDAL treatment for up to 3 years showed no adverse effects on growth and sexual maturation. No differences were observed between risperidone and placebo groups in measurements of sexual maturation, using the Tanner scale, and no adverse events suggestive of delayed pubertal maturation were reported. The mean change in height after 1 year of treatment with risperidone was within the expected growth range in this population.

Experience of risperidone treatment in children with schizophrenia aged less than 15 years is lacking. Experience is lacking in children with conduct and other disruptive behaviour disorders aged less than 5 years. Experience is lacking in children with autism aged less than 5 years. However, in a toxicity study with juvenile rats, increased pup mortality and a delay in physical development was observed. In a 40-week study with juvenile dogs, sexual maturation was delayed. Long bone growth was not affected at a dose similar to the maximum human dose in adolescents (6 mg/day); effects were observed at a dose 4-fold (on an AUC basis) or 7-fold (on a mg/m<sup>2</sup> basis) the maximum human dose in adolescents.

## **ADVERSE EFFECTS**

### **Clinical Trial Data**

The safety of RISPERDAL was evaluated from a clinical trial database consisting of 9803 patients exposed to one or more doses of RISPERDAL for the treatment of various psychiatric disorders in adults, elderly patients with dementia, and pediatrics. Of these 9803 patients, 2687 were patients who received RISPERDAL while participating in double-blind, placebo-controlled trials. The conditions and duration of treatment with RISPERDAL varied greatly and included (in overlapping categories) double-blind, fixed- and flexible-dose, placebo- or active-controlled studies and open-label phases of studies, inpatients and outpatients, and short-term (up to 12 weeks) and longer-term (up to 3 years) exposures.

The majority of all adverse reactions were mild to moderate in severity.

### ***Double-Blind, Placebo-Controlled Data – Adult Patients***

Adverse drug reactions (ADRs) reported by  $\geq 1\%$  of RISPERDAL-treated adult patients in nine 3- to 8-week double-blind, placebo-controlled trials are shown in Table 1.

<b>System/Organ Class</b> Adverse Reaction	<b>RISPERDAL</b> <b><math>\leq 8</math> mg/day</b> <b>(N=853)</b>	<b>RISPERDAL</b> <b><math>&gt;8</math>-16 mg/day</b> <b>(N=198)</b>	<b>PLACEBO</b> <b>(N=687)</b>
	%	%	%
<b>Infections and Infestations</b>			
Nasopharyngitis	2.1	4.0	1.7
Upper respiratory tract infection	1.5	2.5	1.5
Sinusitis	0.7	1.5	0.6
Urinary tract infection	0.5	2.5	0.1

<b>Blood and Lymphatic System Disorders</b>			
Anaemia	0.1	1.0	0.1
<b>Immune System Disorders</b>			
Hypersensitivity	0.1	1.0	0.1
<b>Psychiatric Disorders</b>			
Insomnia	16.2	25.3	13.2
Anxiety	7.7	11.1	4.4
Nervousness	0.5	1.0	0.1
<b>Nervous System Disorders</b>			
Parkinsonism*	19.3	17.2	7.9
Akathisia*	9.8	10.1	2.7
Somnolence	6.8	1.5	2.0
Dizziness	6.3	3.5	3.9
Sedation	4.6	3.0	1.3
Tremor*	4.2	2.5	2.5
Dystonia*	3.8	3.5	1.0
Lethargy	2.6	0	1.3
Dizziness postural	1.2	0	0.1
Dyskinesia*	1.2	2.0	0.9
Syncope	0.4	1.0	0
<b>Eye Disorders</b>			
Vision blurred	2.1	1.0	0.7
<b>Ear and Labyrinth Disorders</b>			
Ear pain	0.1	1.0	0.3
<b>Cardiac Disorders</b>			
Tachycardia	1.1	2.5	0.1
<b>Vascular Disorders</b>			
Orthostatic hypotension	1.3	0.5	0.1
Hypotension	0.2	1.0	0.3
<b>Respiratory, Thoracic and Mediastinal Disorders</b>			
Nasal congestion	2.0	6.1	1.3
Dyspnoea	0.8	2.0	0
Epistaxis	0.5	1.5	0.1
Sinus congestion	0.5	1.0	0.6
<b>Gastrointestinal Disorders</b>			
Nausea	6.4	4.0	2.6
Constipation	4.6	9.1	3.6
Dyspepsia	4.3	6.1	2.6
Vomiting	3.9	4.5	3.8
Diarrhoea	2.3	0.5	1.9
Salivary hypersecretion	2.3	1.0	0.4
Dry mouth	2.1	0	1.0
Abdominal discomfort	1.5	1.0	0.9
Abdominal pain	1.1	0.5	0.7
Stomach discomfort	1.1	1.0	0.6
Abdominal pain upper	0.7	1.0	0.1
<b>Skin and Subcutaneous Tissue Disorders</b>			
Rash	0.8	3.5	0.9
Dry skin	0.5	2.5	0.3
Dandruff	0.2	1.0	0
Seborrhoeic dermatitis	0.2	1.0	0
Hyperkeratosis	0	1.0	0.3
<b>Musculoskeletal and Connective Tissue Disorders</b>			
Back pain	2.5	1.0	1.6

Arthralgia	1.5	2.5	0.6
Pain in extremity	1.2	1.0	2.2
<b>Renal and Urinary Disorders</b>			
Urinary incontinence	0.2	1.0	0.3
<b>Reproductive System and Breast Disorders</b>			
Ejaculation failure	0.4	1.0	0
<b>General Disorders</b>			
Fatigue	2.3	1.0	1.0
Asthenia	1.3	0.5	0.6
Pyrexia	1.3	1.0	0.7
Chest pain	0.8	1.5	0.4
<b>Investigations</b>			
Blood creatine phosphokinase increased	0.4	1.5	0.1
Heart rate increased	0.2	1.5	0.1

\* Parkinsonism includes extrapyramidal disorder, musculoskeletal stiffness, Parkinsonism, cogwheel rigidity, akinesia, bradykinesia, hypokinesia, masked facies, muscle rigidity, and Parkinson's disease. Akathisia includes akathisia and restlessness. Dystonia includes dystonia, muscle spasms, muscle contractions involuntary, muscle contracture, oculogyration, tongue paralysis. Tremor includes tremor and Parkinsonian rest tremor. Dyskinesia includes dyskinesia, muscle twitching, chorea, and choreoathetosis.

### **Double-Blind, Placebo-Controlled Data – Elderly Patients with Dementia**

Adverse drug reactions (ADRs) reported by  $\geq 1\%$  of RISPERDAL-treated elderly patients with dementia in six 4- to 12-week double-blind, placebo-controlled trials are shown in Table 2. Table 2 includes only those ADRs that are either not listed in Table 1 or those ADRs that occurred at  $\geq 2$  times the frequency of the ADRs listed in Table 1.

<b>Table 2.</b> Adverse Drug Reactions (ADRs) Reported by $\geq 1\%$ of RISPERDAL-Treated Elderly Patients with Dementia in Double-Blind Placebo-Controlled Studies: ADRs Not Listed in Table 1 or Reported at $\geq 2$ Times the Frequency of ADRs Listed in Table 1.		
<b>System/Organ Class</b> Adverse Reaction	<b>RISPERDAL</b> <b>(N=1009)</b> %	<b>PLACEBO</b> <b>(N=712)</b> %
<b>Infections and Infestations</b>		
Urinary tract infection	12.9	10.3
Pneumonia	3.1	2.4
Cellulitis	1.1	1.3
<b>Metabolism and Nutrition Disorders</b>		
Decreased appetite	2.3	1.4
<b>Psychiatric Disorders</b>		
Confusional state	2.7	0.1
<b>Nervous System Disorders</b>		
Lethargy	7.6	2.2
Transient ischaemic attack	1.6	0.6
Depressed level of consciousness	1.3	0.3
Drooling	1.3	0
Cerebrovascular accident	1.1	0.4
<b>Eye Disorders</b>		
Conjunctivitis	2.7	1.1
<b>Vascular Disorders</b>		

<b>Table 2.</b> Adverse Drug Reactions (ADRs) Reported by $\geq 1\%$ of RISPERDAL-Treated Elderly Patients with Dementia in Double-Blind Placebo-Controlled Studies: ADRs Not Listed in Table 1 or Reported at $\geq 2$ Times the Frequency of ADRs Listed in Table 1.		
<b>System/Organ Class</b> Adverse Reaction	<b>RISPERDAL</b> <b>(N=1009)</b> %	<b>PLACEBO</b> <b>(N=712)</b> %
Hypotension	2.2	1.4
<b>Respiratory, Thoracic and Mediastinal Disorders</b>		
Cough	4.6	3.1
Rhinorrhoea	1.5	0.8
<b>Gastrointestinal Disorders</b>		
Dysphagia	1.5	1.3
Faecaloma	1.1	0.4
<b>Skin and Subcutaneous Tissue Disorders</b>		
Erythema	4.0	4.6
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Posture abnormal	1.8	0.8
Joint swelling	1.5	0.3
<b>General Disorders</b>		
Oedema peripheral	7.7	3.9
Pyrexia	4.0	1.8
Gait disturbance	3.5	1.5
Pitting oedema	1.5	0.3
<b>Investigations</b>		
Body temperature increased	2.6	0.8

### ***Double-Blind, Placebo-Controlled Data – Pediatric Patients***

Adverse drug reactions (ADRs) reported by  $\geq 1\%$  of RISPERDAL-treated pediatric patients in eight 3- to 8-week double-blind, placebo-controlled trials are shown in Table 3. Table 3 includes only those ADRs that are either not listed in Table 1 or those ADRs that occurred at  $\geq 2$  times the frequency of the ADRs listed in Table 1.

**Table 3.** Adverse Drug Reactions (ADRs) Reported by  $\geq 1\%$  of RISPERDAL-Treated Pediatric Patients in Double-Blind Placebo-Controlled Studies: ADRs Not Listed in Table 1 or Reported at  $\geq 2$  Times the Frequency of ADRs Listed in Table 1.

<b>System/Organ Class</b> Adverse Reaction	<b>RISPERDAL</b> $\leq 3$ mg/day (N=344) %	<b>RISPERDAL</b> >3-6 mg/day (N=95) %	<b>PLACEBO</b> (N=349) %
<b>Infections and Infestations</b>			
Upper respiratory tract infection	5.2	2.1	3.4
Rhinitis	3.5	1.1	3.2
Influenza	1.7	0	1.7
<b>Metabolism and Nutrition Disorders</b>			
Increased appetite	17.2	3.2	7.2
<b>Psychiatric Disorders</b>			
Middle insomnia	1.7	0	0.9
Listless	0.9	1.1	0
<b>Nervous System Disorders</b>			
Somnolence	26.5	15.8	7.7
Headache	22.4	21.1	14.9
Sedation	20.1	14.7	4.0
Dizziness	8.1	13.7	2.3
Tremor	6.1	8.4	1.1
Drooling	4.9	2.1	1.1
Dysarthria	1.5	1.1	0
Disturbance in attention	0.9	1.1	0.6
Balance disorder	0.9	1.1	0
Hypersomnia	0.6	1.1	0.9
<b>Cardiac Disorders</b>			
Palpitations	0.6	2.1	0
<b>Respiratory, Thoracic and Mediastinal Disorders</b>			
Cough	8.7	3.2	6.6
Rhinorrhoea	4.9	2.1	3.4
Epistaxis	3.8	4.2	1.7
Pharyngolaryngeal pain	3.8	2.1	1.7
Pulmonary congestion	0.3	1.1	0.3
<b>Gastrointestinal Disorders</b>			
Vomiting	13.7	8.4	9.2
Abdominal pain upper	8.4	6.3	4.6
Diarrhoea	6.7	2.1	6.0
Salivary hypersecretion	3.5	6.3	0.9
Stomach discomfort	2.9	0	1.4
Abdominal pain	2.3	2.1	0.6
<b>Skin and Subcutaneous Tissue Disorders</b>			
Pruritus	1.2	0	0
Acne	0.9	1.1	0
<b>Musculoskeletal and Connective Tissue Disorders</b>			
Myalgia	1.2	1.1	0.9
Neck pain	0.3	1.1	0.3
<b>Renal and Urinary Disorders</b>			
Enuresis	6.4	1.1	5.2
Urinary incontinence	2.0	0	1.4
Pollakiuria	1.5	1.1	0.3
<b>Reproductive System and Breast Disorders</b>			
Galactorrhea	0.6	2.1	0
<b>General Disorders</b>			
Fatigue	19.2	18.9	4.9
Pyrexia	8.4	3.2	6.3
Feeling abnormal	1.2	0	0
Sluggishness	0.9	1.1	0

Chest discomfort	0.3	1.1	0
<b>Investigations</b>			
Weight increased	4.9	2.1	0.9
Blood prolactin increased	3.8	0	0.3

### Other Clinical Trial Data

Paliperidone is the active metabolite of risperidone, therefore the adverse reaction profiles of these compounds (including both the oral and injectable formulations) are relevant to one another. This subsection includes additional ADRs reported with risperidone and/or paliperidone in clinical trials.

ADRs reported with risperidone and/or paliperidone by  $\geq 1\%$  of RISPERDAL-treated subjects in a pooled dataset of 23 double-blind, placebo-controlled pivotal studies (9 in adults, 6 in elderly patients with dementia, and 8 in paediatric patients) are shown in Table 4a.

<p><b>Table 4a.</b> ADRs Reported with Risperidone and/or Paliperidone by <math>\geq 1\%</math> of RISPERDAL - treated Subjects in a Pooled Dataset of the 23 Double-blind, Placebo-controlled Pivotal Studies- 9 in Adults, 6 in Elderly Patients with Dementia, and 8 in Paediatric patients (The Terms within each System Organ Class are Sorted Alphabetically)</p>
<p><b>System/Organ Class</b> Adverse Reaction <b>Psychiatric disorders</b> Agitation, Insomnia* <b>Nervous System Disorders</b> Akathisia*, Dyskinesia*, Dystonia*, Parkinsonism* <b>Vascular disorders</b> Hypertension <b>Musculoskeletal and Connective Tissue Disorders</b> Musculoskeletal pain <b>General disorders and administration site conditions</b> Gait abnormal, Oedema*, Pain <b>Injury, poisoning and procedural complications</b> Fall</p>

\* **Insomnia includes:** initial insomnia, middle insomnia; **Akathisia includes:** hyperkinesia, restless legs syndrome, restlessness; **Dyskinesia includes:** athetosis, chorea, choreoathetosis, movement disorder, muscle twitching, myoclonus; **Dystonia includes:** blepharospasm, cervical spasm, emprosthotonus, facial spasm, hypertonia, laryngospasm, muscle contractions involuntary, myotonia, oculogyration, opisthotonus, oropharyngeal spasm, pleurothotonus, risus sardonicus, tetany, tongue paralysis, tongue spasm, torticollis, trismus; **Parkinsonism includes:** akinesia, bradykinesia, cogwheel rigidity, drooling, extrapyramidal symptoms, glabellar reflex abnormal, muscle rigidity, muscle tightness, musculoskeletal stiffness; **Oedema includes:** generalised oedema, oedema peripheral, pitting oedema.

ADRs reported with risperidone and/or paliperidone by  $< 1\%$  of RISPERDAL-treated subjects in a pooled dataset of 23 double-blind, placebo-controlled pivotal studies (9 in adults, 6 in elderly patients with dementia, and 8 in paediatric patients) are shown in Table 4b.

<p><b>Table 4b.</b> ADRs Reported with Risperidone and/or Paliperidone by <math>&lt; 1\%</math> of RISPERDAL-treated Subjects in a Pooled Dataset of 23 Double-blind, Placebo-controlled Pivotal Studies -9 in Adults, 6 in Elderly Patients with Dementia, and 8 in Paediatric patients. (The Terms within each System Organ Class are Sorted Alphabetically).</p>
<p><b>System/Organ Class</b> Adverse Reaction <b>Infections and infestations</b> Acarodermatitis, Bronchitis, Cystitis, Ear infection, Eye infection, Infection, Localised infection, Onychomycosis, Respiratory tract infection, Tonsillitis, Viral infection</p>

**Blood and lymphatic system disorders**

Eosinophil count increased, Haematocrit decreased, Neutropenia, White blood cell count decreased

**Endocrine disorders**

Glucose urine present, Hyperprolactinaemia

**Metabolism and nutrition disorders**

Anorexia, Blood cholesterol increased, Blood triglycerides increased, Hyperglycaemia, Polydipsia, Weight decreased

**Psychiatric disorders**

Blunted affect, Depression, Libido decreased, Nightmare, Sleep disorder

**Nervous system disorders**

Cerebrovascular disorder, Convulsion\*, Coordination abnormal, Diabetic coma, Hypoaesthesia, Loss of consciousness, Paraesthesia, Psychomotor hyperactivity, Tardive dyskinesia, Unresponsive to stimuli

**Eye disorders**

Dry eye, Eye rolling, Eyelid margin crusting, Glaucoma, Lacrimation increased, Ocular hyperaemia

**Ear and labyrinth disorders**

Tinnitus, Vertigo

**Cardiac disorders**

Atrioventricular block, Bradycardia, Conduction disorder, Electrocardiogram abnormal, Electrocardiogram QT prolonged, Sinus arrhythmia

**Vascular disorders**

Flushing

**Respiratory, thoracic and mediastinal disorders**

Dysphonia, Hyperventilation, Pneumonia aspiration, Rales, Respiratory disorder, Respiratory tract congestion, Wheezing

**Gastrointestinal disorders**

Cheilitis, Faecal incontinence, Flatulence, Gastroenteritis, Swollen tongue, Toothache

**Hepatobiliary disorders**

Gamma-glutamyltransferase increased, Hepatic enzyme increased, Transaminases increased

**Skin and subcutaneous tissue disorders**

Eczema, Skin discolouration, Skin disorder, Skin lesion

**Musculoskeletal and connective tissue disorders**

Joint stiffness, Muscular weakness, Rhabdomyolysis

**Renal and urinary disorders**

Dysuria

**Reproductive system and breast disorders**

Amenorrhoea, Breast discharge, Ejaculation disorder, Erectile dysfunction, Gynaecomastia, Menstrual disorder\*, Sexual dysfunction, Vaginal discharge

**System/Organ Class**

Adverse Reaction

**General disorders and administration site conditions**

Body temperature decreased, Chills, Discomfort, Drug withdrawal syndrome, Face oedema, Malaise, Peripheral coldness, Thirst

**Injury, poisoning and procedural complications**

Procedural pain

ADRs reported with risperidone and/or paliperidone in other clinical trials but not reported by RISPERDAL-treated subjects in a pooled dataset of 23 double-blind, placebo-controlled pivotal studies are shown in Table 4c.

**Table 4c.** ADRs Reported with Risperidone and/or Paliperidone in Other Clinical Trials but Not Reported by RISPERDAL-treated Subjects in a Pooled Dataset of 23 Double-blind, Placebo-controlled Pivotal Studies. (The Terms within each System Organ Class are Sorted Alphabetically)

**System/Organ Class**

Adverse Reaction

**Immune system disorders**

Anaphylactic reaction

**Metabolism and nutrition disorders**

Hyperinsulinaemia

**Psychiatric disorders**

Anorgasmia

**Nervous system disorders**

Head titubation, Neuroleptic malignant syndrome

**Eye disorders**

Eye movement disorder, Photophobia

**Cardiac disorders**

Postural orthostatic tachycardia syndrome

**Gastrointestinal disorders**

Intestinal obstruction

**Skin and subcutaneous tissue disorders**

Drug eruption, Urticaria

**Reproductive system and breast disorders**

Breast discomfort, Breast engorgement, Breast enlargement, Menstruation delayed

**General disorders and administration site conditions**

Induration

**Class effects**

QT prolongation, ventricular arrhythmias (ventricular fibrillation, ventricular tachycardia), sudden unexplained death, cardiac arrest and Torsade de pointes may occur with antipsychotics.

**Postmarketing Data**

Adverse events first identified as ADRs during postmarketing experience with risperidone and/or paliperidone are included in Tables 5. The frequencies are provided according to the following convention:

Very common	≥1/10
Common	≥1/100 to <1/10
Uncommon	≥1/1,000 to <1/100
Rare	≥1/10,000 to <1/1,000
Very rare	<1/10,000, including isolated reports

In Table 5, ADRs are presented by frequency category based on spontaneous reporting rate.

**Table 5.** Adverse Drug Reactions Identified During Postmarketing Experience with Risperidone and/or Paliperidone by Frequency Category Estimated from Spontaneous Reporting Rates with Risperidone

**Blood and Lymphatic Disorders***Very rare* Agranulocytosis, Thrombocytopenia**Endocrine Disorders***Very rare* Inappropriate antidiuretic hormone secretion**Metabolism and Nutrition Disorders***Very rare* Diabetes mellitus, Diabetic ketoacidosis, Hypoglycaemia, Water intoxication**Psychiatric Disorders***Very rare* Mania**Nervous System Disorders***Very rare* Dysgeusia**Eye Disorders***Very rare* Floppy iris syndrome (intraoperative)**Cardiac Disorders***Very rare* Atrial fibrillation**Vascular Disorders***Very rare* Deep vein thrombosis, Pulmonary embolism

**Respiratory, Thoracic, and Mediastinal Disorders**

*Very rare* Sleep apnoea syndrome

**Gastrointestinal Disorders**

*Very rare* Pancreatitis, ileus

**Hepatobiliary Disorders**

*Very rare* Jaundice

**Skin and Subcutaneous Tissue Disorders**

*Very rare* Alopecia, Angioedema

**Renal and Urinary Disorders**

*Very rare* Urinary retention

**Pregnancy, Puerperium and Perinatal Conditions**

*Very rare* Drug withdrawal syndrome neonatal

**Reproductive System and Breast Disorders**

*Very rare* Priapism

**General Disorders**

*Very rare* Hypothermia

There have also been reports of benign pituitary adenoma that were temporally related, but not necessarily causally related, to risperidone therapy.

## INTERACTIONS

### Pharmacodynamic-related Interactions

#### *Centrally-acting Drugs and Alcohol*

Given the primary CNS effects of risperidone, it should be used with caution in combination with other centrally-acting medicines or alcohol.

#### *Levodopa and Dopamine Agonists*

Risperidone may antagonise the effects of levodopa and other dopamine agonists.

#### *Drugs with Hypotensive Effects*

Clinically significant hypotension has been observed postmarketing with concomitant use of risperidone and antihypertensive treatment.

#### *Drugs Known to Prolong the QT interval*

As with other antipsychotics, caution should be exercised when RISPERDAL is prescribed in combination with other medicines thought to prolong the QT interval or medicines known to cause electrolyte imbalance.

### Pharmacokinetic-related Interactions

Food does not affect the absorption of RISPERDAL.

Risperidone is mainly metabolised through CYP2D6, and to a lesser extent through CYP3A4. Both risperidone and its active metabolite 9-hydroxyrisperidone are substrates of P-glycoprotein (P-gp). Substances that modify CYP2D6 activity, or substances strongly inhibiting or inducing CYP3A4 and/or P-gp activity, may influence the pharmacokinetics of the risperidone active antipsychotic fraction.

#### *Strong CYP2D6 Inhibitors*

Co-administration of RISPERDAL with a strong CYP2D6 inhibitor may increase the plasma concentrations of risperidone, but less so of the active antipsychotic fraction. Higher doses of a

strong CYP2D6 inhibitor may elevate concentrations of the risperidone active antipsychotic fraction (e.g., paroxetine, see below). When concomitant paroxetine or another strong CYP2D6 inhibitor, especially at higher doses, is initiated or discontinued, the physician should re-evaluate the dosing of RISPERDAL.

#### *CYP3A4 and/or P-gp Inhibitors*

Coadministration of RISPERDAL with a strong CYP3A4 and/or P-gp inhibitor may substantially elevate plasma concentrations of the risperidone active antipsychotic fraction. When concomitant itraconazole or another strong CYP3A4 and/or P-gp inhibitor is initiated or discontinued, the physician should re-evaluate the dosing of RISPERDAL.

#### *CYP3A4 and/or P-gp Inducers*

Co-administration of RISPERDAL with a strong CYP3A4 and/or P-gp inducer may decrease the plasma concentrations of the risperidone active antipsychotic fraction. When concomitant carbamazepine or another strong CYP3A4 and/or P-gp inducer is initiated or discontinued, the physician should re-evaluate the dosing of RISPERDAL.

#### *Highly Protein-bound Drugs*

When RISPERDAL is taken together with highly protein-bound drugs, there is no clinically relevant displacement of either drug from the plasma proteins.

When using concomitant medication, the corresponding label should be consulted for information on the route of metabolism and the possible need to adjust dosages.

#### *Paediatric Population*

Interaction studies have only been performed in adults. The relevance of the results from these studies in pediatric patients is unknown.

#### *Examples*

Examples of drugs that may potentially interact or that were shown not to interact with risperidone are listed below:

##### Antibacterials:

- Erythromycin, a moderate CYP3A4 inhibitor, does not change the pharmacokinetics of risperidone and the active antipsychotic fraction.
- Rifampicin, a strong CYP3A4 inducer and a P-gp inducer, decreased the plasma concentrations of the active antipsychotic fraction.

##### Anticholinesterases:

- Donepezil and galantamine, both CYP2D6 and CYP3A4 substrates, do not show a clinically relevant effect on the pharmacokinetics of risperidone and the active antipsychotic fraction.

##### Antiepileptics:

- Carbamazepine, a strong CYP3A4 inducer and a P-gp inducer, has been shown to decrease the plasma concentrations of the active antipsychotic fraction of risperidone
- Topiramate modestly reduced the bioavailability of risperidone, but not that of the active antipsychotic fraction. Therefore, this interaction is unlikely to be of clinical significance.
- Risperidone does not show a clinically relevant effect on the pharmacokinetics of valproate or topiramate.

#### Antifungals:

- Itraconazole, a strong CYP3A4 inhibitor and a P-gp inhibitor, at a dosage of 200 mg/day increased the plasma concentrations of the active antipsychotic fraction by about 70%, at risperidone doses of 2 to 8 mg/day.
- Ketoconazole, a strong CYP3A4 inhibitor and a P-gp inhibitor, at a dosage of 200 mg/day increased the plasma concentrations of risperidone and decreased the plasma concentrations of 9-hydroxyrisperidone.

#### Antipsychotics:

- Phenothiazines may increase the plasma concentrations of risperidone but not those of the active antipsychotic fraction.
- Aripiprazole, a CYP2D6 and CYP3A4 substrate: Risperidone tablets or injections did not affect the pharmacokinetics of the sum of aripiprazole and its active metabolite, dehydroaripiprazole.

#### Antivirals:

- Protease inhibitors: No formal study data are available; however, since ritonavir is a strong CYP3A4 inhibitor and a weak CYP2D6 inhibitor, ritonavir and ritonavir-boosted protease inhibitors potentially raise concentrations of the risperidone active antipsychotic fraction.

#### Beta Blockers:

- Some beta-blockers may increase the plasma concentrations of risperidone but not those of the active antipsychotic fraction.

#### Calcium Channel Blockers:

- Verapamil, a moderate inhibitor of CYP3A4 and an inhibitor of P-gp, increases the plasma concentration of risperidone and the active antipsychotic fraction.

#### Digitalis Glycosides:

- Risperidone does not show a clinically relevant effect on the pharmacokinetics of digoxin.

#### Diuretics:

- Furosemide: See section Warnings and Precautions regarding increased mortality in elderly patients with dementia concomitantly receiving furosemide and oral RISPERDAL.

#### Gastrointestinal Drugs:

- H2-receptor antagonists: Cimetidine and ranitidine, both weak inhibitors of CYP2D6 and CYP3A4, increased the bioavailability of risperidone, but only marginally that of the active antipsychotic fraction.

#### Lithium:

- Risperidone does not show a clinically relevant effect on the pharmacokinetics of lithium.

#### Sodium Channel Blockers:

- Quinidine may increase the plasma concentrations of risperidone but not those of the active antipsychotic fraction.

#### SSRIs and Tricyclic Antidepressants:

- Fluoxetine, a strong CYP2D6 inhibitor, increases the plasma concentration of risperidone but less so of the active antipsychotic fraction.
- Paroxetine, a strong CYP2D6 inhibitor, increases the plasma concentrations of risperidone, but, at dosages up to 20 mg/day, less so of the active antipsychotic fraction.

However, higher doses of paroxetine may elevate concentrations of the risperidone active antipsychotic fraction.

- Sertraline, a weak inhibitor of CYP2D6, and fluvoxamine, a weak inhibitor of CYP3A4, at dosages up to 100 mg/day are not associated with clinically significant changes in concentrations of the risperidone active antipsychotic fraction. However, doses higher than 100 mg/day of sertraline or fluvoxamine may elevate concentrations of the risperidone active antipsychotic fraction.
- Tricyclic antidepressants may increase the plasma concentrations of risperidone but not those of the active antipsychotic fraction. Amitriptyline does not affect the pharmacokinetics of risperidone or the active antipsychotic fraction (see **Pharmacokinetics**).

## OVERDOSAGE

### Symptoms

In general, reported signs and symptoms have been those resulting from an exaggeration of the known pharmacological effects of RISPERDAL. These include drowsiness and sedation, tachycardia and hypotension, and extrapyramidal symptoms. In overdose, QT prolongation and convulsions have been reported. Torsade de pointes has been reported in association with combined overdose of oral RISPERDAL and paroxetine.

### Treatment

Establish and maintain a clear airway and ensure adequate oxygenation and ventilation. Gastric lavage (after intubation, if the patient is unconscious) and administration of activated charcoal together with a laxative should be considered. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias.

There is no specific antidote to RISPERDAL. Therefore appropriate supportive measures should be instituted. Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluids and/or sympathomimetic agents. In case of severe extrapyramidal symptoms, anticholinergic medication should be administered. Close medical supervision and monitoring should continue until the patient recovers.

As strategies for the management of overdose are continually evolving, it is advisable to contact the Poisons Information Centre to determine the latest recommendations for the management of an overdose.

## PHARMACEUTICAL PRECAUTIONS

### Shelf Life

RISPERDAL 0.5 mg Tablets: 2 years when stored below 30°C.

RISPERDAL 1 mg, 2 mg, 3 mg and 4 mg Tablets: 3 years when stored below 25°C in a dry place and protected from light.

RISPERDAL Quicklet 0.5 mg, 1 mg, 2 mg, 3 mg and 4 mg tablets: 2 years when stored below 30°C.

RISPERDAL Oral Solution should be stored below 30°C and should be protected from freezing. Keep out of reach of children.

## MEDICINE CLASSIFICATION

Prescription Medicine

## **PACKAGE QUANTITIES**

RISPERDAL 0.5 mg Tablets - blisters in a carton of 20.

RISPERDAL 1 mg, 2 mg, 3 mg and 4 mg Tablets - blisters in a carton of 60.

RISPERDAL Quicklet 0.5 mg, 1 mg, 2 mg, 3 mg and 4 mg Tablets-blisters in a carton of 28.

RISPERDAL Oral Solution 1 mg/mL - bottle in 30 mL with a pipette of 3 mL, calibrated in mg and mL. Minimum volume is 0.25 mL; maximum volume is 3 mL.

Not all presentations may be marketed.

## **NAME AND ADDRESS**

Janssen-Cilag (New Zealand) Ltd  
Auckland, NEW ZEALAND  
Telephone: 0800 800 806

## **DATE OF PREPARATION:**

11 July 2014