

**TYLENOL<sup>®</sup>  
with Codeine**

**C<sub>III</sub>**

**(acetaminophen and codeine phosphate) tablets**

**WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; HEPATOTOXICITY; CYTOCHROME P450 2D6 INTERACTION; DEATH RELATED TO ULTRA-RAPID METABOLISM OF CODEINE TO MORPHINE; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS**

**Addiction, Abuse and Misuse**

TYLENOL<sup>®</sup> with Codeine exposes patients and other users to the risks of opioid addiction, abuse and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing TYLENOL<sup>®</sup> with Codeine, and monitor all patients regularly for the development of these behaviors and conditions [see WARNINGS].

**Life-Threatening Respiratory Depression**

Serious, life-threatening, or fatal respiratory depression may occur with use of TYLENOL<sup>®</sup> with Codeine. Monitor for respiratory depression, especially during initiation of TYLENOL<sup>®</sup> with Codeine or following a dose increase [see WARNINGS].

**Accidental Ingestion**

Accidental ingestion of TYLENOL<sup>®</sup> with Codeine, especially by children, can result in a fatal overdose of TYLENOL<sup>®</sup> with Codeine [see WARNINGS].

**Neonatal Opioid Withdrawal Syndrome**

Prolonged use of TYLENOL<sup>®</sup> with Codeine during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see WARNINGS].

**Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Respiratory depression and death have occurred in children who received codeine following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine due to a CYP2D6 polymorphism [see WARNINGS, PRECAUTIONS; Information for Patients/Caregivers, Nursing Mothers].

### **Hepatotoxicity**

Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4,000 milligrams per day, and often involve more than one acetaminophen-containing product [see WARNINGS].

### **Interactions with Drugs Affecting Cytochrome P450 Isoenzymes**

The effects of concomitant use or discontinuation of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with codeine are complex. Use of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with **TYLENOL<sup>®</sup>** with Codeine requires careful consideration of the effects on the parent drug, codeine, and the active metabolite, morphine.

- **Cytochrome P450 3A4 Interaction**

The concomitant use of **TYLENOL<sup>®</sup>** with Codeine with all cytochrome P450 3A4 inhibitors or discontinuation of a cytochrome P450 3A4 inducer may result in an increase in codeine plasma concentrations with subsequently greater metabolism by cytochrome P450 2D6, resulting in greater morphine levels, which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression.

The concomitant use of **TYLENOL<sup>®</sup>** with Codeine with all cytochrome P450 3A4 inducers or discontinuation of a cytochrome P450 3A4 inhibitor may result in lower codeine levels, greater norcodeine levels, and less metabolism via 2D6 with resultant lower morphine levels. This may be associated with a decrease in efficacy, and in some patients, may result in signs and symptoms of opioid withdrawal.

Follow patients receiving **TYLENOL<sup>®</sup>** with Codeine and any CYP3A4 inhibitor or inducer for signs and symptoms that may reflect opioid toxicity and opioid withdrawal when **TYLENOL<sup>®</sup>** with Codeine is used in conjunction with inhibitors and inducers of CYP3A4 [see WARNINGS, PRECAUTIONS; Drug Interactions].

- **Cytochrome P450 2D6 Interaction**

The concomitant use of **TYLENOL<sup>®</sup>** with Codeine with all cytochrome P450 2D6 inhibitors may result in an increase in codeine plasma concentrations and a decrease in the plasma concentration of the active metabolite, morphine, which could result in an analgesic efficacy reduction or symptoms of opioid withdrawal.

The discontinuation of a cytochrome P450 2D6 inhibitor may result in a decrease in codeine plasma concentrations and an increase in the plasma concentration of the active metabolite, morphine, which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression.

Follow patients receiving **TYLENOL<sup>®</sup>** with Codeine and any CYP2D6 inhibitor for signs and symptoms that may reflect opioid toxicity and opioid withdrawal when **TYLENOL<sup>®</sup>** with Codeine are used in conjunction with inhibitors of CYP2D6 [see WARNINGS, PRECAUTIONS; Drug Interactions].

**Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**


Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death [see WARNINGS , PRECAUTIONS; Drug Interactions].

- Reserve concomitant prescribing of **TYLENOL<sup>®</sup>** with Codeine and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

## DESCRIPTION

TYLENOL<sup>®</sup> with Codeine is supplied in tablet form for oral administration.


Acetaminophen, 4'-hydroxyacetanilide, a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



$C_8 H_9 NO_2$

M.W. 151.16

Codeine phosphate, 7,8-didehydro-4, 5 $\alpha$ -epoxy-3-methoxy-17-methylmorphinan-6 $\alpha$ -ol phosphate (1:1) (salt) hemihydrate, a white crystalline powder, is a narcotic analgesic and antitussive. It has the following structural formula:



$C_{18}H_{21}NO_3 \cdot H_3PO_4 \cdot 1/2 H_2O$

M.W. 406.37

Each Tylenol with Codeine No. 3 tablet, USP (300 mg/30 mg) contains:

Acetaminophen	300 mg
Codeine Phosphate	30 mg

Each Tylenol with Codeine No. 4 tablet, USP (300 mg/60 mg) contains:

Acetaminophen	300 mg
Codeine Phosphate	60 mg

In addition, each tablet contains the following inactive ingredients:

TYLENOL<sup>®</sup> with Codeine No. 3 contains powdered cellulose, magnesium stearate, sodium metabisulfite\*, pregelatinized starch (corn), and modified starch (corn).

TYLENOL<sup>®</sup> with Codeine No. 4 contains powdered cellulose, magnesium stearate, sodium metabisulfite\*, pregelatinized starch (corn), and corn starch.

\* See **WARNINGS**

## **CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Codeine is an opioid agonist relatively selective for the mu-opioid receptor, but with a much weaker affinity than morphine. The analgesic properties of codeine have been speculated to come from its conversion to morphine, although the exact mechanism of analgesic action remains unknown.

The precise mechanism of the analgesic properties of acetaminophen is not established but is thought to involve central actions.

### **Pharmacodynamics**

#### **Effects on the Central Nervous System**

Codeine produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

Codeine causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

#### **Effects on the Gastrointestinal Tract and Other Smooth Muscle**

Codeine causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm, resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, and transient elevations in serum amylase.

### Effects on the Cardiovascular System

Codeine produces peripheral vasodilation which may result in orthostatic hypotension or syncope. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes, sweating, and/or orthostatic hypotension.

### Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotrophic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans [see **ADVERSE REACTIONS**]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see **ADVERSE REACTIONS**].

### Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

### Concentration–Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. The minimum effective analgesic concentration of codeine for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome, and/or the development of analgesic tolerance [see **DOSAGE AND ADMINISTRATION**].

### Concentration–Adverse Reaction Relationships

There is a relationship between increasing codeine plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see **DOSAGE AND ADMINISTRATION**].

### Pharmacokinetics

The behavior of the individual components is described below.

## Codeine

Codeine is rapidly absorbed from the gastrointestinal tract. It is rapidly distributed from the intravascular spaces to the various body tissues, with preferential uptake by parenchymatous organs such as the liver, spleen, and kidney. Codeine crosses the blood-brain barrier and is found in fetal tissue and breast milk. The plasma concentration does not correlate with brain concentration or relief of pain. Codeine is about 7-25% bound to plasma proteins and does not accumulate in body tissues.

About 70 to 80% of the administered dose of codeine is metabolized by conjugation with glucuronic acid to codeine-6-glucuronide (C6G) and via O-demethylation to morphine (about 5 to 10%) and N-demethylation to norcodeine (about 10%) respectively. UDP-glucuronosyltransferase (UGT) 2B7 and 2B4 are the major enzymes mediating glucuronidation of codeine to C6G. Cytochrome P450 2D6 is the major enzyme responsible for conversion of codeine to morphine and P450 3A4 is the major enzyme mediating conversion of codeine to norcodeine. Morphine and norcodeine are further metabolized by conjugation with glucuronic acid. The glucuronide metabolites of morphine are morphine-3-glucuronide (M3G) and morphine-6-glucuronide (M6G). Morphine and M6G are known to have analgesic activity in humans. The analgesic activity of C6G in humans is unknown. Norcodeine and M3G are generally not considered to possess analgesic properties.

The plasma half-life is about 2.9 hours. The elimination of codeine is primarily via the kidneys, and about 90% of an oral dose is excreted by the kidneys within 24 hours of dosing. The urinary secretion products consist of free and glucuronide conjugated codeine (about 70%), free and conjugated norcodeine (about 10%), free and conjugated morphine (about 10%), normorphine (4%), and hydrocodone (1%). The remainder of the dose is excreted in the feces.

At therapeutic doses, the analgesic effect reaches a peak within 2 hours and persists between 4 and 6 hours.

## Acetaminophen

Acetaminophen is rapidly absorbed from the gastrointestinal tract and is distributed throughout most body tissues. A small fraction (10-25%) of acetaminophen is bound to plasma proteins. The plasma half-life is 1.25 to 3 hours, but may be increased by liver damage and following overdose. Elimination of acetaminophen is principally by liver metabolism (conjugation) and subsequent renal excretion of metabolites. Acetaminophen is primarily metabolized in the liver by first-order kinetics and involves three principal separate pathways: conjugation with glucuronide; conjugation with sulfate; and oxidation via the cytochrome, P450-dependent, mixed-

function oxidase enzyme pathway to form a reactive intermediate metabolite, which conjugates with glutathione and is then further metabolized to form cysteine and mercapturic acid conjugates. The principal cytochrome P450 isoenzyme involved appears to be CYP2E1, with CYP1A2 and CYP3A4 as additional pathways. Approximately 85% of an oral dose appears in the urine within 24 hours of administration, most as the glucuronide conjugate, with small amounts of other conjugates and unchanged drug.

See **OVERDOSAGE** for toxicity information.

## **INDICATIONS AND USAGE**

TYLENOL<sup>®</sup> with Codeine (acetaminophen and codeine phosphate) is indicated for the management of mild to moderate pain where treatment with an opioid is appropriate and for which alternative treatments are inadequate.

### Limitations of Use

Because of the risks of addiction, abuse, and misuse, with opioids, even at recommended doses [see **WARNINGS**], reserve TYLENOL<sup>®</sup> with Codeine for use in patients for whom alternative treatment options [e.g., non-opioid analgesics]

- Have not provided adequate analgesia, or are not expected to provide adequate analgesia,
- Have not been tolerated, or are not expected to be tolerated.

## **CONTRAINDICATIONS**

TYLENOL<sup>®</sup> with Codeine is contraindicated in:

- Patients with significant respiratory depression [see **WARNINGS**].
- Patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [see **WARNINGS**].
- Postoperative pain management in children who have undergone tonsillectomy and/or adenoidectomy [see **WARNINGS**].
- Patients with known or suspected gastrointestinal obstruction, including paralytic ileus [see **WARNINGS**].
- Patients with hypersensitivity to codeine, acetaminophen, or any of the formulation excipients (e.g., anaphylaxis) [see **WARNINGS**].

- Concurrent use of monoamine oxidase inhibitors (MAOIs) or use of MAOIs within the last 14 days [see **WARNINGS**].

## **WARNINGS**

### **Addiction, Abuse, and Misuse**

TYLENOL<sup>®</sup> with Codeine contains codeine, a Schedule III controlled substance. As an opioid, TYLENOL<sup>®</sup> with Codeine exposes users to the risks of addiction, abuse, and misuse [see **DRUG ABUSE and DEPENDENCE**].

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed TYLENOL<sup>®</sup> with Codeine. Addiction can occur at recommended dosages and if the drug is misused or abused.

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing TYLENOL<sup>®</sup> with Codeine, and monitor all patients receiving TYLENOL<sup>®</sup> with Codeine for the development of these behaviors and conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed opioids such as TYLENOL<sup>®</sup> with Codeine, but use in such patients necessitates intensive counseling about the risks and proper use of TYLENOL<sup>®</sup> with Codeine along with intensive monitoring for signs of addiction, abuse, and misuse.

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when prescribing or dispensing TYLENOL<sup>®</sup> with Codeine. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug [see **PRECAUTIONS; Information for Patients/Caregivers**]. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

### **Life-Threatening Respiratory Depression**

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status

[see **OVERDOSAGE**]. Carbon dioxide (CO<sub>2</sub>) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of TYLENOL<sup>®</sup> with Codeine, the risk is greatest during the initiation of therapy or following a dosage increase. Monitor patients closely for respiratory depression, especially within the first 24-72 hours of initiating therapy with and following dosage increases of TYLENOL<sup>®</sup> with Codeine.

To reduce the risk of respiratory depression, proper dosing and titration of TYLENOL<sup>®</sup> with Codeine are essential [see **DOSAGE AND ADMINISTRATION**]. Overestimating the TYLENOL<sup>®</sup> with Codeine dosage when converting patients from another opioid product can result in a fatal overdose with the first dose.

Accidental ingestion of TYLENOL<sup>®</sup> with Codeine, especially by children, can result in respiratory depression and death due to an overdose of codeine.

### **Neonatal Opioid Withdrawal Syndrome**

Prolonged use of TYLENOL<sup>®</sup> with Codeine during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for a prolonged period of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see **PRECAUTIONS, Information for Patients/Caregivers, Pregnancy**].

### **Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Codeine-containing products are contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy [see **CONTRAINDICATIONS**].

Respiratory depression and death have occurred in children who received codeine in the postoperative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for cytochrome P450 isoenzyme 2D6 or high morphine concentrations). Deaths have also occurred in nursing infants who were exposed to high levels of morphine in breast milk because their mothers were ultra-rapid metabolizers of codeine [see **PRECAUTIONS; Information for Patients/Caregivers, Nursing Mothers**].

Some individuals may be ultra-rapid metabolizers because of a specific CYP2D6 genotype (gene duplications denoted as \*1/\*1xN or \*1/\*2xN). The prevalence of this CYP2D6 phenotype varies widely and has been estimated at 0.5 to 1% in Chinese and Japanese, 0.5 to 1% in Hispanics, 1 to 10% in Caucasians, 3% in African Americans, and 16 to 28% in North Africans, Ethiopians, and Arabs. Data are not available for other ethnic groups. These individuals convert codeine into its active metabolite, morphine, more rapidly and completely than other people. This rapid conversion results in higher than expected serum morphine levels. Even at labeled dosage regimens, individuals who are ultra-rapid metabolizers may have life-threatening or fatal respiratory depression or experience signs of overdose (such as extreme sleepiness, confusion, or shallow breathing) [see **OVERDOSAGE**].

Children with obstructive sleep apnea who are treated with codeine for post-tonsillectomy and/or adenoidectomy pain may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine.

### **Risks of Interactions with Drugs Affecting Cytochrome P450 Isoenzymes**

The effects of concomitant use or discontinuation of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with codeine are complex. Use of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with **TYLENOL<sup>®</sup>** with Codeine requires careful consideration of the effects on the parent drug, codeine, and the active metabolite, morphine.

#### **Cytochrome P450 3A4 Interaction**

The concomitant use of **TYLENOL<sup>®</sup>** with Codeine with all cytochrome P450 3A4 inhibitors, such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir) or discontinuation of a cytochrome P450 3A4 inducer such as rifampin, carbamazepine, and phenytoin, may result in an increase in codeine plasma concentrations with subsequently greater metabolism by cytochrome P450 2D6, resulting in greater morphine levels, which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression.

The concomitant use of **TYLENOL<sup>®</sup>** with Codeine with all cytochrome P450 3A4 inducers or discontinuation of a cytochrome P450 3A4 inhibitor may result in lower codeine levels, greater norcodeine levels, and less metabolism via 2D6 with resultant lower morphine levels. This may be associated with a decrease in efficacy, and in some patients, may result in signs and symptoms of opioid withdrawal.

Follow patients receiving TYLENOL<sup>®</sup> with Codeine and any CYP3A4 inhibitor or inducer for signs and symptoms that may reflect opioid toxicity and opioid withdrawal when TYLENOL<sup>®</sup> with Codeine are used in conjunction with inhibitors and inducers of CYP3A4 [see **WARNINGS, PRECAUTIONS, Drug Interactions**].

### **Risks of Concomitant Use or Discontinuation of Cytochrome P450 2D6 Inhibitors**

The concomitant use of TYLENOL<sup>®</sup> with Codeine with all cytochrome P450 2D6 inhibitors (e.g., amiodarone, quinidine) may result in an increase in codeine plasma concentrations and a decrease in active metabolite morphine plasma concentration which could result in an analgesic efficacy reduction or symptoms of opioid withdrawal.

Discontinuation of a concomitantly used cytochrome P450 2D6 inhibitor may result in a decrease in codeine plasma concentration and an increase in active metabolite morphine plasma concentration which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression.

Follow patients receiving TYLENOL<sup>®</sup> with Codeine and any CYP2D6 inhibitor for signs and symptoms that may reflect opioid toxicity and opioid withdrawal when TYLENOL<sup>®</sup> with Codeine are used in conjunction with inhibitors of CYP2D6 [see **PRECAUTIONS, Drug Interactions**].

### **Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of TYLENOL<sup>®</sup> with Codeine with benzodiazepines and/or other CNS depressants (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol). Because of these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics [see **PRECAUTIONS; Drug Interactions**].

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid

analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression and sedation.

Advise both patients and caregivers about the risks of respiratory depression and sedation when TYLENOL<sup>®</sup> with Codeine is used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS depressants including alcohol and illicit drugs [see **PRECAUTIONS; Drug Interactions, Information for Patients/Caregivers**].

### **Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or Elderly, Cachectic, or Debilitated Patients**

The use of TYLENOL<sup>®</sup> with Codeine in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated.

#### **Patients with Chronic Pulmonary Disease**

TYLENOL<sup>®</sup> with Codeine-treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of TYLENOL<sup>®</sup> with Codeine [see **WARNINGS; Life-Threatening Respiratory Depression**].

#### **Elderly, Cachectic, or Debilitated Patients**

Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics, including clearance, compared to younger, healthier patients [see **WARNINGS; Respiratory Depression**].

Monitor such patients closely, particularly when initiating and titrating TYLENOL<sup>®</sup> with Codeine and when TYLENOL<sup>®</sup> with Codeine is given concomitantly with other drugs that depress respiration [see **WARNINGS; Life-Threatening Respiratory**

**Depression**]. Alternatively, consider the use of non-opioid analgesics in these patients.

### **Interaction with Monoamine Oxidase Inhibitors**

Monoamine oxidase inhibitors (MAOIs) may potentiate the effects of morphine, codeine's active metabolite, including respiratory depression, coma, and confusion. TYLENOL<sup>®</sup> with Codeine should not be used in patients taking MAOIs or within 14 days of stopping such treatment.

### **Adrenal Insufficiency**

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than 1 month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

### **Severe Hypotension**

TYLENOL<sup>®</sup> with Codeine may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [see **PRECAUTIONS; Drug Interactions**]. Monitor these patients for signs of hypotension after initiating or titrating the dosage of TYLENOL<sup>®</sup> with Codeine. In patients with circulatory shock TYLENOL<sup>®</sup> with Codeine may cause vasodilatation that can further reduce cardiac output and blood pressure. Avoid the use of TYLENOL<sup>®</sup> with Codeine with circulatory shock.

### **Hepatotoxicity**

Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4,000 milligrams per day, and often involve more than one acetaminophen-containing product. The excessive intake of acetaminophen may be intentional to cause self-harm or unintentional as patients

attempt to obtain more pain relief or unknowingly take other acetaminophen-containing products.

The risk of acute liver failure is higher in individuals with underlying liver disease and in individuals who ingest alcohol while taking acetaminophen.

Instruct patients to look for acetaminophen or APAP on package labels and not to use more than one product that contains acetaminophen. Instruct patients to seek medical attention immediately upon ingestion of more than 4,000 milligrams of acetaminophen per day, even if they feel well.

### **Serious Skin Reactions**

Rarely, acetaminophen may cause serious skin reactions such as acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. Patients should be informed about the signs of serious skin reactions, and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

### **Hypersensitivity/Anaphylaxis**

There have been post-marketing reports of hypersensitivity and anaphylaxis associated with the use of acetaminophen. Clinical signs included swelling of the face, mouth, and throat, respiratory distress, urticaria, rash, pruritus, and vomiting. There were infrequent reports of life-threatening anaphylaxis requiring emergency medical attention. Instruct patients to discontinue TYLENOL<sup>®</sup> with Codeine immediately and seek medical care if they experience these symptoms. Do not prescribe TYLENOL<sup>®</sup> with Codeine for patients with acetaminophen allergy [see **PRECAUTIONS; Information for Patients/Caregivers**].

### **Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness**

In patients who may be susceptible to the intracranial effects of CO<sub>2</sub> retention (e.g., those with evidence of increased intracranial pressure or brain tumors), TYLENOL<sup>®</sup> with Codeine may reduce respiratory drive, and the resultant CO<sub>2</sub> retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when initiating therapy with TYLENOL<sup>®</sup> with Codeine.

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of TYLENOL<sup>®</sup> with Codeine in patients with impaired consciousness or coma.

### **Risks of Use in Patients with Gastrointestinal Conditions**

TYLENOL<sup>®</sup> with Codeine is contraindicated in patients with gastrointestinal obstruction, including paralytic ileus.

The administration of TYLENOL<sup>®</sup> with Codeine or other opioids may obscure the diagnosis or clinical course in patients with acute abdominal conditions.

TYLENOL<sup>®</sup> with Codeine may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

### **Sulfite Sensitivity**

TYLENOL<sup>®</sup> with Codeine contains sodium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

### **Increased Risk of Seizures in Patients with Seizure Disorders**

The codeine in TYLENOL<sup>®</sup> with Codeine may increase the frequency of seizures in patients with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during TYLENOL<sup>®</sup> with Codeine therapy.

### **Withdrawal**

Avoid the use of mixed agonist/antagonist (e.g, pentazocine, nalbuphine, and butorphanol) or partial agonist (e.g., buprenorphine) analgesics in patients who are receiving a full opioid agonist analgesic, including TYLENOL<sup>®</sup> with Codeine. In these patients, mixed agonist/antagonist and partial analgesics may reduce the analgesic effect and/or precipitate withdrawal symptoms.

When discontinuing TYLENOL<sup>®</sup> with Codeine, gradually taper the dosage [see **DOSAGE AND ADMINISTRATION**]. Do not abruptly discontinue TYLENOL<sup>®</sup> with Codeine [see **DRUG ABUSE AND DEPENDENCE**].

## **PRECAUTIONS**

### **Risks of Driving and Operating Machinery**

TYLENOL<sup>®</sup> with Codeine may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to

the effects of TYLENOL<sup>®</sup> with Codeine and know how they will react to the medication [see **PRECAUTIONS; Information for Patients/Caregivers**].

### **Information for Patients/Caregivers**

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

#### **Addiction, Abuse and Misuse**

Inform patients that the use of TYLENOL<sup>®</sup> with Codeine, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [see **WARNINGS**]. Instruct patients not to share TYLENOL<sup>®</sup> with Codeine with others and to take steps to protect TYLENOL<sup>®</sup> with Codeine from theft or misuse.

#### **Life-Threatening Respiratory Depression**

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting TYLENOL<sup>®</sup> with Codeine or when the dosage is increased, and that it can occur even at recommended dosages [see **WARNINGS**]. Advise patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop.

#### **Accidental Ingestion**

Inform patients that accidental ingestion, especially by children, may result in respiratory depression or death [see **WARNINGS**]. Instruct patients to take steps to store TYLENOL<sup>®</sup> with Codeine securely. Advise patients to properly dispose of TYLENOL<sup>®</sup> with Codeine in accordance with local state guidelines and/or regulations.

#### **Interactions with Benzodiazepines and Other CNS Depressants**

Inform patients and caregivers that potentially fatal additive effects may occur if TYLENOL<sup>®</sup> with Codeine is used with benzodiazepines or other CNS depressants, including alcohol, and not to use these drugs concomitantly unless supervised by a healthcare provider [see **WARNINGS, PRECAUTIONS; Drug Interactions**].

#### **Serotonin Syndrome**

Inform patients that opioids could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Warn patients of the symptoms and signs of serotonin syndrome and to seek medical attention right away if symptoms develop.

Instruct patients to inform their healthcare provider if they are taking, or plan to take serotonergic medications [see **PRECAUTIONS; Drug Interactions**].

### MAOI Interaction

Inform patients not to take TYLENOL<sup>®</sup> with Codeine while using any drugs that inhibit monoamine oxidase. Patients should not start MAOIs while taking TYLENOL<sup>®</sup> with Codeine [see **WARNINGS, PRECAUTIONS; Drug Interactions**].

### Adrenal Insufficiency

Inform patients that opioids could cause adrenal insufficiency, a potentially life-threatening condition. Adrenal insufficiency may present with non-specific symptoms and signs such as nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. Advise patients to seek medical attention if they experience a constellation of these symptoms [see **WARNINGS**].

### Important Administration Instructions

Instruct patients how to properly take TYLENOL<sup>®</sup> with Codeine [see **DOSAGE AND ADMINISTRATION**].

### Maximum Daily Dose of Acetaminophen

Inform patients not to take more than 4,000 milligrams of acetaminophen per day. Advise patients to call their healthcare provider if they have taken more than the recommended dose.

### Hypotension

Inform patients that TYLENOL<sup>®</sup> with Codeine may cause orthostatic hypotension and syncope. Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious consequences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position) [see **WARNINGS; Hypotension**].

### Anaphylaxis

Inform patients that anaphylaxis has been reported with ingredients contained in TYLENOL<sup>®</sup> with Codeine. Advise patients how to recognize such a reaction, and if they develop signs of allergy such as a rash or difficulty breathing to stop taking TYLENOL<sup>®</sup> with Codeine and seek medical attention. [see **CONTRAINDICATIONS, ADVERSE REACTIONS**].

### Pregnancy

#### *Neonatal Opioid Withdrawal Syndrome*

Inform female patients of reproductive potential that prolonged use of TYLENOL<sup>®</sup> with Codeine during pregnancy can result in neonatal opioid withdrawal syndrome,

which may be life-threatening if not recognized and treated [see **WARNINGS, PRECAUTIONS; Pregnancy**].

#### *Embryo-Fetal Toxicity*

Inform female patients of reproductive potential that TYLENOL<sup>®</sup> with Codeine can cause fetal harm and to inform the prescriber of a known or suspected pregnancy [see **PRECAUTIONS; Pregnancy**].

#### **Lactation**

Advise patients that nursing mothers taking codeine can have higher morphine levels in their breast milk if they are ultra-rapid metabolizers. These higher levels of morphine in breast milk may lead to life-threatening or fatal side effects in nursing babies. Advise nursing mothers to watch for signs of morphine toxicity in their infants which includes increased sleepiness (more than usual), difficulty breastfeeding, breathing difficulties, or limpness. Instruct nursing mothers to talk to the baby's doctor immediately if they notice these signs and, if they cannot reach the doctor right away, to take the baby to an emergency room or call 911 (or local emergency services) [see **PRECAUTIONS; Nursing Mothers**].

#### **Infertility**

Inform patients that chronic use of opioids may cause reduced fertility. It is not known whether these effects on fertility are reversible.

#### **Driving or Operating Heavy Machinery**

Inform patients that TYLENOL<sup>®</sup> with Codeine may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery and to avoid such tasks while taking this product, until they know how they will react to the medication.

#### **Ultra-Rapid Metabolism of Codeine**

Advise patients that some people have a genetic variation that results in codeine changing into morphine more rapidly and completely than other people. Most people are unaware of whether they are an ultra-rapid codeine metabolizer or not. These higher-than-normal levels of morphine in the blood may lead to life-threatening or fatal respiratory depression or signs of overdose such as extreme sleepiness, confusion, or shallow breathing. Children with this genetic variation who were prescribed codeine after tonsillectomy and/or adenoidectomy for obstructive sleep apnea may be at greatest risk based on reports of several deaths in this population due to respiratory depression. Codeine-containing products are contraindicated in all children who undergo tonsillectomy and/or adenoidectomy. Advise caregivers of

children receiving codeine-containing products for other reasons to monitor for signs of respiratory depression.

### **Disposal of Unused TYLENOL<sup>®</sup> with Codeine**

Advise patients to properly dispose of TYLENOL<sup>®</sup> with Codeine tablets. Advise patients to throw the drug in the household trash following these steps:

1. Remove them from their original containers and mix them with an undesirable substance, such as used coffee grounds or kitty litter (this makes the drug less appealing to children and pets, and unrecognizable to people who may intentionally go through the trash seeking drugs).
2. Place the mixture in a sealable bag, empty can, or other container to prevent the drug from leaking or breaking out of a garbage bag, or dispose of unused tablets in accordance with local state guidelines and/or regulations.

### **Drug Interactions**

#### **CYP2D6 Inhibitors**

Codeine is metabolized by CYP2D6 to form morphine. The concomitant use of TYLENOL<sup>®</sup> with Codeine and CYP2D6 inhibitors (e.g., paroxetine, fluoxetine, bupropion, quinidine) can increase the plasma concentration of codeine, but decreases the plasma concentration of active metabolite morphine, particularly when an inhibitor is added after a stable dose of TYLENOL<sup>®</sup> with Codeine is achieved.

If concomitant use is necessary, consider dosage adjustment of TYLENOL<sup>®</sup> with Codeine until stable drug effects are achieved.

After stopping a CYP2D6 inhibitor, as the effects of the inhibitor decline, the codeine plasma concentration will decrease but the morphine plasma concentration will increase. If a CYP2D6 inhibitor is discontinued, consider adjusting the TYLENOL<sup>®</sup> with Codeine dosage until stable drug effects are achieved.

If concomitant use is necessary or if a CYP2D6 inhibitor is discontinued after concomitant use, monitor patients closely at frequent intervals. If signs and symptoms of respiratory depression or sedation occur, consider reducing the TYLENOL<sup>®</sup> with Codeine dosage until stable drug effects are achieved.

#### **CYP3A4 Inhibitors**

The concomitant use of TYLENOL<sup>®</sup> with Codeine and CYP3A4 inhibitors such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g. ketoconazole), and protease inhibitors (e.g., ritonavir), can increase the plasma concentration of codeine, resulting in increased or prolonged opioid effects,

particularly when an inhibitor is added after a stable dose of TYLENOL<sup>®</sup> with Codeine is achieved. If concomitant use is necessary, consider dosage reduction of TYLENOL<sup>®</sup> with Codeine until stable drug effects are achieved. Monitor patients for respiratory depression and sedation at frequent intervals.

After stopping a CYP3A4 inhibitor, as the effects of the inhibitor decline, the codeine plasma concentration will decrease resulting in decreased opioid efficacy or a withdrawal syndrome in patients who had developed physical dependence to codeine. If a CYP3A4 inhibitor is discontinued, consider increasing the TYLENOL<sup>®</sup> with Codeine dosage until stable drug effects are achieved. Monitor for signs of opioid withdrawal.

### CYP3A4 Inducers

The concomitant use of TYLENOL<sup>®</sup> with Codeine and CYP3A4 inducers (e.g., rifampin, carbamazepine, phenytoin) can decrease the plasma concentration of codeine, resulting in decreased efficacy or onset of a withdrawal syndrome in patients who have developed physical dependence to codeine. If concomitant use is necessary, consider increasing the TYLENOL<sup>®</sup> with Codeine dosage until stable drug effects are achieved. Monitor for signs of opioid withdrawal.

After stopping a CYP3A4 inducer, as the effects of the inducer decline, the codeine plasma concentration will increase, which could increase or prolong both the therapeutic effects and adverse reactions, and may cause serious respiratory depression. If a CYP3A4 inducer is discontinued, consider a TYLENOL<sup>®</sup> with Codeine dosage reduction and monitor for signs of respiratory depression.

### Benzodiazepines and Other Central Nervous System (CNS) Depressants

Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants, including alcohol, other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics and other opioids, can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death.

Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Follow patients closely for signs of respiratory depression and sedation [see **WARNINGS**].

### Serotonergic Drugs

The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system has resulted in serotonin syndrome. Examples of these drugs

include, selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT<sub>3</sub> receptor antagonists, drugs that effect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), and monoamine oxidase (MAO) inhibitors (used to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue) [see **PRECAUTIONS; Information for Patients**].

If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation and dose adjustment. Discontinue **TYLENOL<sup>®</sup>** with Codeine immediately if serotonin syndrome is suspected.

#### **Monoamine Oxidase Inhibitors (MAOIs)**

The concomitant use of opioids and MAOIs, such as phenelzine, tranylcypromine, linezolid, may manifest as serotonin syndrome or opioid toxicity.

Advise patients taking **TYLENOL<sup>®</sup>** with Codeine not to use MAOIs or within 14 days of stopping such treatment. If urgent use of an opioid is necessary, use test doses and frequent titration of small doses of other opioids (such as oxycodone, hydrocodone, oxymorphone, hydrocodone, or buprenorphine) to treat pain while closely monitoring blood pressure and signs and symptoms of CNS and respiratory depression.

#### **Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics**

The concomitant use of opioids with other opioid analgesics, such as butorphanol, nalbuphine, pentazocine, may reduce the analgesic effect of **TYLENOL<sup>®</sup>** with Codeine and/or precipitate withdrawal symptoms.

Advise patient to avoid concomitant use of these drugs.

#### **Muscle Relaxants**

**TYLENOL<sup>®</sup>** with Codeine may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.

If concomitant use is warranted, monitor patients for signs of respiratory depression that may be greater than otherwise expected and decrease the dosage of acetaminophen and codeine phosphate tablet and/or the muscle relaxant as necessary.

#### **Diuretics**

Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone.

If concomitant use is warranted, monitor patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.

### **Anticholinergic Drugs**

The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.

If concomitant use is warranted, monitor patients for signs of urinary retention or reduced gastric motility when TYLENOL<sup>®</sup> with Codeine is used concomitantly with anticholinergic drugs.

### **Drug/Laboratory Test Interactions**

Codeine may increase serum amylase levels.

Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenesis**

Long-term studies to evaluate the carcinogenic potential of the combination of codeine and acetaminophen have not been conducted.

Two-year carcinogenicity studies have been conducted in F344/N rats and B6C3F1 mice. There was no evidence of carcinogenicity in male and female rats, respectively, at dietary doses up to 70 and 80 mg/kg/day of codeine sulfate (approximately 2 times the maximum recommended daily dose of 360 mg/day for adults on a mg/m<sup>2</sup> basis) for two years. Similarly there was no evidence of carcinogenicity activity in male and female mice at dietary doses up to 400 mg/kg/day of codeine sulfate (approximately 5 times the maximum recommended daily dose of 360 mg/day for adults on a mg/m<sup>2</sup> basis) for two years.

Long-term studies in mice and rats have been completed by the National Toxicology Program to evaluate the carcinogenic potential of acetaminophen. In 2-year feeding studies, F344/N rats and B6C3F1 mice were fed a diet containing acetaminophen up to 6000 ppm. Female rats demonstrated equivocal evidence of carcinogenic activity based on increased incidences of mononuclear cell leukemia at 0.8 times the maximum human daily dose (MHDD) of 4 grams/day, based on a body surface area comparison. In contrast, there was no evidence of carcinogenic activity in male rats that received up to 0.7 times or mice at up to 1.2-1.4 times the MHDD, based on a body surface area comparison.

## Mutagenesis

Codeine sulfate was not mutagenic in the *in vitro* bacterial reverse mutation assay or clastogenic in the *in vitro* Chinese hamster ovary cell chromosome aberration assay.

In the published literature, acetaminophen has been reported to be clastogenic when administered at 1500 mg/kg/day to the rat model (3.6-times the MHDD, based on a body surface area comparison). In contrast, no clastogenicity was noted at a dose of 750 mg/kg/day (1.8-times the MHDD, based on a body surface area comparison), suggesting a threshold effect.

## Impairment of Fertility

No nonclinical fertility studies have been conducted with codeine or the combination of codeine and acetaminophen.

In studies conducted by the National Toxicology Program, fertility assessments with acetaminophen have been completed in Swiss CD-1 mice via a continuous breeding study. There were no effects on fertility parameters in mice consuming up to 1.7 times the MHDD of acetaminophen, based on a body surface area comparison. Although there was no effect on sperm motility or sperm density in the epididymis, there was a significant increase in the percentage of abnormal sperm in mice consuming 1.78 times the MHDD (based on a body surface comparison) and there was a reduction in the number of mating pairs producing a fifth litter at this dose, suggesting the potential for cumulative toxicity with chronic administration of acetaminophen near the upper limit of daily dosing.

Published studies in rodents report that oral acetaminophen treatment of male animals at doses that are 1.2 times the MHDD and greater (based on a body surface comparison) result in decreased testicular weights, reduced spermatogenesis, reduced fertility, and reduced implantation sites in females given the same doses. These effects appear to increase with the duration of treatment. The clinical significance of these findings is not known.

## Infertility

Chronic use of opioids may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [see **ADVERSE REACTIONS**].

## **Pregnancy**

Teratogenic Effects: Pregnancy Category C

### *Codeine*

A study in rats and rabbits reported no teratogenic effect of codeine administered during the period of organogenesis in doses ranging from 5 to 120 mg/kg. In the rat, doses at the 120 mg/kg level, in the toxic range for the adult animal, were associated with an increase in embryo resorption at the time of implantation. In another study a single 100 mg/kg subcutaneous dose of codeine administered to pregnant mice reportedly resulted in delayed ossification in the offspring.

There are no adequate and well-controlled studies in pregnant women. TYLENOL<sup>®</sup> with Codeine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## **Nonteratogenic Effects**

### *Fetal/Neonatal Adverse Reactions*

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [see **WARNINGS**].

## **Labor or Delivery**

Opioids cross the placenta and may produce respiratory depression and psychophysiologic effects in neonates. An opioid antagonist, such as naloxone, must be available for reversal of opioid-induced respiratory depression in the neonate. TYLENOL<sup>®</sup> with Codeine is not recommended for use in pregnant women during or immediately prior to labor, when other analgesic techniques are more appropriate. Opioid analgesics, including TYLENOL<sup>®</sup> with Codeine, and can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Narcotic analgesics should be avoided during labor if delivery of a premature infant is anticipated. If the mother has received narcotic analgesics during labor, newborn infants should be observed closely for signs of respiratory depression. Resuscitation may be required [see **OVERDOSAGE**]. The effect of codeine, if any, on the later growth, development, and functional maturation of the child is unknown.

### **Nursing Mothers**

Codeine is secreted into human milk. In women with normal codeine metabolism (normal CYP2D6 activity), the amount of codeine secreted into human milk is low and dose-dependent. However, some women are ultra-rapid metabolizers of codeine. These women achieve higher-than-expected serum levels of codeine's active metabolite, morphine, leading to higher-than-expected levels of morphine in breast milk and potentially dangerously high serum morphine levels in their breastfed infants. Therefore, maternal use of codeine can potentially lead to serious adverse reactions, including death, in nursing infants.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for **TYLENOL<sup>®</sup> with Codeine** and any potential adverse effects on the breastfed infant from **TYLENOL<sup>®</sup> with Codeine** or from the underlying maternal condition.

The risk of infant exposure to codeine and morphine through breast milk should be weighed against the benefits of breastfeeding for both the mother and baby. Caution should be exercised when codeine is administered to a nursing woman. If a codeine-containing product is selected, the lowest dose should be prescribed for the shortest period of time to achieve the desired clinical effect. Infants exposed to codeine phosphate through breast milk should be monitored for excess sedation and respiratory depression. Mothers using codeine should be informed about when to seek immediate medical care and how to identify the signs and symptoms of neonatal toxicity, such as drowsiness or sedation, difficulty breastfeeding, breathing difficulties, and decreased tone, in their baby. Nursing mothers who are ultra-rapid metabolizers may also experience overdose symptoms such as extreme sleepiness, confusion, or shallow breathing. Prescribers should closely monitor mother-infant pairs and notify treating pediatricians about the use of codeine during breastfeeding [see **WARNINGS; Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**].

Acetaminophen is excreted in breast milk in small amounts, but the significance of its effect on nursing infants is not known. Because of the potential for serious adverse reactions in nursing infants from acetaminophen, a decision should be made whether

to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Infants exposed to TYLENOL<sup>®</sup> with Codeine through breast milk should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

### **Pediatric Use**

Respiratory depression and death have occurred in children with obstructive sleep apnea who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for cytochrome P450 isoenzyme CYP2D6 or high morphine concentrations). These children may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine-containing products are contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy [see **CONTRAINDICATIONS, WARNINGS**].

### **Geriatric Use**

Elderly patients (aged 65 years or older) may have increased sensitivity to TYLENOL<sup>®</sup> with Codeine. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids, and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of TYLENOL<sup>®</sup> with Codeine slowly in geriatric patients and monitor closely for signs of central nervous system depression [see **WARNINGS**].

These drugs are known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

The following adverse reactions have been identified during post-approval use of TYLENOL<sup>®</sup> with Codeine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Addiction, Abuse, and Misuse [see **WARNINGS**]
- Life-Threatening Respiratory Depression [see **WARNINGS**]
- Neonatal Opioid Withdrawal Syndrome [see **WARNINGS**]
- Ultra-rapid Metabolizers of Codeine [see **WARNINGS**]
- Interactions with CNS Depressants [see **WARNINGS**]
- Severe Hypotension [see **WARNINGS**]
- Gastrointestinal Adverse Reactions [see **WARNINGS**]
- Seizures [see **WARNINGS**]
- Withdrawal [see **WARNINGS**]

Serious adverse reactions associated with codeine are respiratory depression and, to a lesser degree, circulatory depression, respiratory arrest, shock, and cardiac arrest.

The most frequently observed adverse reactions with codeine administration include drowsiness, lightheadedness, dizziness, sedation, shortness of breath, nausea, vomiting, sweating, and constipation.

Other adverse reactions include allergic reactions, euphoria, dysphoria, abdominal pain, pruritus, rash, thrombocytopenia, and agranulocytosis.

Other less frequently observed adverse reactions expected from opioid analgesics, including TYLENOL<sup>®</sup> with Codeine:

*Cardiovascular system:* faintness, flushing, hypotension, palpitations, syncope.

*Digestive System:* abdominal cramps, anorexia, diarrhea, dry mouth, gastrointestinal distress, pancreatitis.

*Nervous system:* anxiety, drowsiness, fatigue, headache, insomnia, nervousness, shakiness, somnolence, vertigo, visual disturbances, weakness.

*Skin and Appendages:* rash, sweating, urticarial.

- Serotonin syndrome: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.
- Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use.
- Anaphylaxis: Anaphylaxis has been reported with ingredients contained in TYLENOL<sup>®</sup> with Codeine.
- Androgen deficiency: Cases of androgen deficiency have occurred with chronic use of opioids [see **CLINICAL PHARMACOLOGY**].

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

TYLENOL<sup>®</sup> with Codeine contains codeine, a Schedule III controlled substance.

### **Abuse**

TYLENOL<sup>®</sup> with Codeine contains codeine, a substance with a high potential for abuse similar to other opioids, including fentanyl, hydrocodone, hydromorphone, methadone, morphine, oxycodone, oxymorphone, and tapentadol. TYLENOL<sup>®</sup> with Codeine can be abused and is subject to misuse, addiction, and criminal diversion [see **WARNINGS**].

All patients treated with opioids require careful monitoring for signs of abuse and addiction, because use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its rewarding psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful, or potentially harmful, consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

“Drug-seeking” behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated “loss” of prescriptions, tampering with prescriptions and reluctance to provide prior medical records or contact information for other treating health care providers. “Doctor shopping” (visiting multiple prescribers to obtain additional prescriptions) is common among drug abusers and people suffering from untreated addiction. Preoccupation

with achieving adequate pain relief can be appropriate behavior in a patient with poor pain control.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Healthcare providers should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

TYLENOL<sup>®</sup> with Codeine, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

#### Risks Specific to Abuse of Acetaminophen and Codeine Phosphate Tablets

Acetaminophen and Codeine Phosphate Tablets is for oral use only. Abuse of Acetaminophen and Codeine Phosphate Tablets poses a risk of overdose and death. The risk is increased with concurrent use of Acetaminophen and Codeine Phosphate Tablets with alcohol and other central nervous system depressants.

#### **Dependence**

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

Physical dependence results in withdrawal symptoms after abrupt discontinuation or a significant dosage reduction of a drug. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

TYLENOL<sup>®</sup> with Codeine should not be abruptly discontinued [see **DOSAGE AND ADMINISTRATION**]. If TYLENOL<sup>®</sup> with Codeine is abruptly discontinued in a physically dependent patient, a withdrawal syndrome may occur. Some or all of the

following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including: irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see **PRECAUTIONS; Pregnancy**].

## **OVERDOSAGE**

Following an acute overdosage, toxicity may result from codeine or acetaminophen.

### **Clinical Presentation**

#### **Codeine**

Acute overdosage with codeine can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations.

#### **Acetaminophen**

Dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect of acetaminophen overdose. Renal tubular necrosis, hypoglycemic coma, and coagulation defects may also occur.

Early symptoms following a potentially hepatotoxic overdose may include; anorexia, nausea, vomiting, diaphoresis, pallor and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion.

### **Treatment of Overdose**

#### **Codeine**

In case of codeine overdose, priorities are the reestablishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or serious arrhythmias will require advanced life-support measures.

The opioid antagonists, naloxone or nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. For clinically significant respiratory or

circulatory depression secondary to TYLENOL<sup>®</sup> with Codeine overdose, administer an opioid antagonist. Only administer opioid antagonists in the presence of clinically significant respiratory, circulatory and/or central nervous system depression secondary to codeine overdose. In patients who are physically dependent on any opioid agonist including TYLENOL<sup>®</sup> with Codeine, an abrupt or complete reversal of opioid effects may precipitate an acute withdrawal syndrome. The severity of the withdrawal syndrome produced will depend on the degree of physical dependence and the dose of the antagonist administered. Please see the prescribing information for the specific opioid antagonist for details of their proper use.

Because the duration of opioid reversal is expected to be less than the duration of action of TYLENOL<sup>®</sup> with Codeine, carefully monitor the patient until spontaneous respiration is reliably reestablished. If the response to an opioid antagonist is suboptimal or only brief in nature, administer additional antagonist as directed by the product's prescribing information.

### Acetaminophen

Gastric decontamination with activated charcoal should be administered just prior to N-acetylcysteine (NAC) to decrease systemic absorption if acetaminophen ingestion is known or suspected to have occurred within a few hours of presentation.

Serum acetaminophen levels should be obtained immediately if the patient presents 4 hours or more after ingestion to assess potential risk of hepatotoxicity; acetaminophen levels drawn less than 4 hours post-ingestion may be misleading. To obtain the best possible outcome, (NAC) should be administered as soon as possible where impending or evolving liver injury is suspected. Intravenous NAC may be administered when circumstances preclude oral administration.

Vigorous supportive therapy is required in severe intoxication. Procedures to limit the continuing absorption of the drug must be readily performed since the hepatic injury is dose-dependent and occurs early in the course of intoxication.

## **DOSAGE AND ADMINISTRATION**

### **Important Dosage and Administration Instructions**

Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [see **WARNINGS**].

Initiate the dosing regimen for each patient individually, taking into account the patient's severity of pain, patient response, prior analgesic treatment experience, and risk factors for addiction, abuse, and misuse [see **WARNINGS**].

Monitor patients closely for respiratory depression, especially within the first 24-72 hours of initiating therapy and following dosage increases with TYLENOL<sup>®</sup> with Codeine and adjust the dosage accordingly [see **WARNINGS**].

### **Initial Dosage**

#### **Initiating Treatment with TYLENOL<sup>®</sup> with Codeine**

Dosage should be adjusted according to severity of pain and response of the patient. However, it should be kept in mind that tolerance to codeine can develop with continued use and that the incidence of untoward effects is dose related. Adult doses of codeine higher than 60 mg are associated with an increased incidence of adverse reactions and are not associated with greater efficacy.

The usual adult dosage is:

TYLENOL<sup>®</sup> with Codeine (codeine 30 mg and acetaminophen 300 mg): Take 1 to 2 tablets every 4 hours as needed for pain.

TYLENOL<sup>®</sup> with Codeine (codeine 60 mg and acetaminophen 300 mg): Take one tablet every 4 hours as needed for pain.

	<b>Single Doses (Range)</b>	<b>Maximum 24-Hour Dose</b>
Codeine Phosphate	30 mg to 60 mg	360 mg
Acetaminophen	300 mg to 1,000 mg	4,000 mg

The prescriber must determine the number of tablets per dose, and the maximum number of tablets per 24 hours, based upon the above dosage guidance. This information should be conveyed in the prescription.

#### **Conversion from Other Opioids to TYLENOL<sup>®</sup> with Codeine**

There is inter-patient variability in the potency of opioid drugs and opioid formulations. Therefore, a conservative approach is advised when determining the total daily dosage of TYLENOL<sup>®</sup> with Codeine. It is safer to underestimate a patient's 24-hour TYLENOL<sup>®</sup> with Codeine dosage than to overestimate the 24-hour TYLENOL<sup>®</sup> with Codeine dosage and manage an adverse reaction due to overdose.

### **Titration and Maintenance of Therapy**

Individually titrate TYLENOL<sup>®</sup> with Codeine to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving TYLENOL<sup>®</sup> with Codeine to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, or misuse [see **WARNINGS**]. Frequent communication is important

among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration.

If the level of pain increases after dosage stabilization, attempt to identify the source of increased pain before increasing the TYLENOL<sup>®</sup> with Codeine dosage. If unacceptable opioid-related adverse reactions are observed, consider reducing the dosage. Adjust the dosage to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

### **Discontinuation of TYLENOL<sup>®</sup> with Codeine**

When a patient who has been taking TYLENOL<sup>®</sup> with Codeine regularly and may be physically dependent no longer requires therapy with TYLENOL<sup>®</sup> with Codeine, taper the dose gradually, by 25% to 50% every 2 to 4 days, while monitoring carefully for signs and symptoms of withdrawal. If the patient develops these signs or symptoms, raise the dose to the previous level and taper more slowly, either by increasing the interval between decreases, decreasing the amount of change in dose, or both. Do not abruptly discontinue TYLENOL<sup>®</sup> with Codeine in a physically-dependent patient [see **WARNINGS, DRUG ABUSE AND DEPENDENCE**].

### **HOW SUPPLIED**

TYLENOL<sup>®</sup> with Codeine (acetaminophen and codeine phosphate) No. 3 Tablets contain acetaminophen 300 mg and codeine phosphate 30 mg. The tablets are white, round, flat-faced, beveled-edged, and imprinted “McNEIL” on one side and “TYLENOL CODEINE” and “3” on the other side.

TYLENOL<sup>®</sup> with Codeine (acetaminophen and codeine phosphate) No. 4 Tablets contain acetaminophen 300 mg and codeine phosphate 60 mg. The tablets are white, round, flat-faced, beveled-edged, and imprinted “McNEIL” on one side and “TYLENOL CODEINE” and “4” on the other side.

TYLENOL<sup>®</sup> with Codeine Tablets are supplied as follows:

TYLENOL<sup>®</sup> with Codeine No. 3 bottle of 100 tablets – NDC 50458-513-60

TYLENOL<sup>®</sup> with Codeine No. 3 bottle of 1000 tablets – NDC 50458-513-80

TYLENOL<sup>®</sup> with Codeine No. 4 bottle of 100 tablets – NDC 50458-515-60

TYLENOL<sup>®</sup> with Codeine No. 4 bottle of 500 tablets – NDC 50458-515-70

Store TYLENOL<sup>®</sup> with Codeine at 20° to 25°C (68° to 77°F). (See USP Controlled Room Temperature.)

Dispense in tight, light-resistant container as defined in the USP.

Manufactured by:  
Janssen Ortho, LLC  
Gurabo, Puerto Rico 00778

Manufactured for:  
Janssen Pharmaceuticals, Inc.  
Titusville, NJ 08560

Revised 01/2017

© 2000 Janssen Pharmaceutical Companies

**Medication Guide**  
**TYLENOL<sup>®</sup> (TAHY-luh-nawl) with Codeine Tablets, CIII**

**TYLENOL<sup>®</sup> with Codeine is:**

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage mild to moderate pain, when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death.

**Important information about TYLENOL<sup>®</sup> with Codeine:**

- **Get emergency help right away if you take too much TYLENOL<sup>®</sup> with Codeine (overdose).** When you first start taking TYLENOL<sup>®</sup> with Codeine, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.
- TYLENOL<sup>®</sup> with Codeine can cause severe drowsiness, breathing problems (respiratory depression), coma and death when taken with benzodiazepines or other medicines that depress consciousness.
- Never give anyone else your TYLENOL<sup>®</sup> with Codeine. They could die from taking it. Store TYLENOL<sup>®</sup> with Codeine away from children and in a safe place to prevent stealing or abuse. Selling or giving away TYLENOL<sup>®</sup> with Codeine is against the law.

**Do not take TYLENOL<sup>®</sup> with Codeine if you have:**

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or narrowing of the stomach or intestines.
- previously had an allergic reaction to codeine or acetaminophen.

**Before taking TYLENOL<sup>®</sup> with Codeine, tell your healthcare provider if you have a history of:**

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

**Tell your healthcare provider if you are:**

- **pregnant or planning to become pregnant.** Prolonged use of TYLENOL<sup>®</sup> with Codeine during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** TYLENOL<sup>®</sup> with Codeine passes into breast milk and may harm your baby.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking TYLENOL<sup>®</sup> with Codeine with certain other medicines can cause serious side effects that could lead to death.

**When taking TYLENOL<sup>®</sup> with Codeine:**

- Do not change your dose. Take TYLENOL<sup>®</sup> with Codeine exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Take your prescribed dose every 4 hours as needed. Do not take more than your prescribed dose. If you miss a dose, take your next dose when needed.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking TYLENOL<sup>®</sup> with Codeine regularly, do not stop taking TYLENOL<sup>®</sup> with Codeine without talking to your healthcare provider.
- After you stop taking TYLENOL<sup>®</sup> with Codeine dispose of any unused tablets in accordance with local state guidelines and/or regulations.

**While taking TYLENOL<sup>®</sup> with Codeine DO NOT:**

- Drive or operate heavy machinery, until you know how TYLENOL<sup>®</sup> with Codeine affects you. TYLENOL<sup>®</sup> with Codeine can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with TYLENOL<sup>®</sup> with Codeine may cause you to overdose and die.

**The possible side effects of TYLENOL<sup>®</sup> with Codeine:**

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

**Get emergency medical help if you have:**

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of TYLENOL<sup>®</sup> with Codeine. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. For more information go to [dailymed.nlm.nih.gov](http://dailymed.nlm.nih.gov).

Manufactured by: Janssen Ortho, LLC, Gurabo, Puerto Rico 00778. Manufactured for: Janssen Pharmaceuticals, Inc., Titusville, NJ 08560. 1-800-526-7736

**This Medication Guide has been approved by the U.S. Food and Drug Administration.**

Issued: December 2016

039822-170210