



Instructions for Use TREMFYA® (guselkumab) Pre-filled Syringe



SINGLE-USE DEVICE

Important

If your doctor decides that you or a caregiver may be able to give your injections of TREMFYA at home, you should receive training on the right way to prepare and inject TREMFYA using the pre-filled syringe before attempting to inject.

Please read this Instructions for Use before using the TREMFYA pre-filled syringe and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your medical condition or your treatment.

The TREMFYA pre-filled syringe is intended for injection under the skin, not into the muscle or vein. After injection, the needle will retract into the body of the device and lock into place.

Storage information


Store in refrigerator at 2° to 8°C. **Do not freeze.**

Keep TREMFYA pre-filled syringe and all medicines out of reach of children.


Do not shake the pre-filled syringe at any time.

Pre-filled safety syringe at-a-glance

Before injection



After injection



You will need these supplies:

- 1 Alcohol swab
- 1 Cotton ball or gauze pad
- 1 Adhesive bandage
- 1 Sharps container (See Step 3)

1. Prepare for your injection



Inspect carton

Remove carton with the pre-filled syringe from the refrigerator.

Keep the pre-filled syringe in the carton and let it sit on a flat surface at room temperature for **at least 30 minutes** before use.

Do not warm any other way.

Check the expiration date ('EXP') on the back panel of the carton.

Do not use if the expiration date has passed.

Do not inject if the perforations on the carton are broken.

Call your doctor or pharmacist for a refill.




Choose injection site

Select from the following areas for your injection:

- **Front of thighs** (recommended)
- Lower abdomen (**do not** use the 5 centimetre area around your belly-button)
- Back of upper arms (if a caregiver is giving you the injection)

Do not inject into skin that is tender, bruised, red, scaly or hard.

Do not inject into areas with scars or stretch marks.




Clean injection site

Wash your hands well with soap and warm water.

Wipe your chosen injection site with an alcohol swab and allow it to dry.

Do not touch, fan or blow on the injection site after you have cleaned it.




Inspect liquid

Take the pre-filled syringe out of the carton.

Check the liquid in the viewing window. It should be clear to slightly yellow and may contain tiny white or clear particles. You may also see one or more air bubbles. This is normal.

Do not inject if the liquid is cloudy or discoloured, or has large particles. Call your doctor or pharmacist for a refill.

2. Inject TREMFYA using pre-filled syringe



Remove needle cover

Hold syringe by the body and pull needle cover straight off.


It is normal to see a drop of liquid.

Inject within 5 minutes of removing the needle cover.

Do not put needle cover back on, as this may damage the needle.

Do not touch needle or let it touch any surface.

Do not use the TREMFYA pre-filled syringe if it is dropped. Call your doctor or pharmacist for a refill.



Position fingers and insert needle


Place your thumb, index and middle fingers **directly under the finger flange**, as shown.

Do not touch plunger or area above finger flange as this may cause the needle safety device to activate.

Use your other hand to pinch skin at the injection site. Position syringe at about a 45 degree angle to the skin.


It is important to pinch enough skin to **inject under the skin** and not into the muscle.

Insert needle with a quick, dart-like motion.




Release pinch and reposition hand

Use your free hand to grasp the body of the syringe.



Press plunger


Place thumb from the opposite hand on the plunger and press the plunger **all the way down until it stops.**



Release pressure from plunger

The safety guard will cover the needle and lock into place, removing the needle from your skin.


3. After your injection



Throw the used pre-filled syringe away

Put your used syringe in a sharps disposal container right away after use.

Make sure you dispose of the bin as instructed by your doctor or nurse when the container is full.



Check injection site

There may be a small amount of blood or liquid at the injection site. Hold pressure over your skin with a cotton ball or gauze pad until any bleeding stops.

Do not rub the injection site.

If needed, cover injection site with a bandage. Your injection is now complete!

Need help?

call your doctor to talk about any questions you may have.

The Janssen Immunology Patient Support Program is available to patients prescribed TREMFYA. **Call 1800 666 845**

© 2022 Janssen Pharmaceutical Companies



Date of preparation: March 2022

Janssen-Cilag Pty Ltd
1-5 Khartoum Road
Macquarie Park NSW 2113





Shared mailbox: RA-JANBE-GRAPHDES@its.jnj.com

IFU TREMFYA
PCC 38413B

Template Ref. Name: Schaffhausen/IFU/3410/1
656x262

Tech. info/Spec:
Market: AU
Mat. ID Code: PCC
File Name: 38413B.indd (CC - PC)
Replaces PCC: 38413A
Font Type: Helvetica World
Font Size: 9 pt

x-height (lowercase) Times NR	Times New Roman compared to our standard fonts				
	Times New Roman	Helvetica World	Antique Olive	Futura	Karbon
0.947 mm	6 pt xxx	5.2 pt xxx	4.3 pt xxx	5.8 pt xxx	6.2 pt xxx
1.104 mm	7 pt xxx	6 pt xxx	5.1 pt xxx	6.7 pt xxx	7.2 pt xxx
1.262 mm	8 pt xxx	6.9 pt xxx	5.8 pt xxx	7.7 pt xxx	8.2 pt xxx
1.42 mm	9 pt xxx	7.7 pt xxx	6.5 pt xxx	8.6 pt xxx	9.2 pt xxx

	Date (Designer)
 Cyan	1. 23-MAY-22 (Jesse)
 Magenta	2.
 Yellow	3.
 Black	4.
	5.
	6.