

**READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE  
PATIENT MEDICATION INFORMATION**

**P<sup>r</sup>PARIET<sup>®</sup>  
rabeprazole sodium tablets**

Read this carefully before you start taking **PARIET<sup>®</sup>** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **PARIET<sup>®</sup>**.

**What is PARIET<sup>®</sup> used for?**

**PARIET<sup>®</sup>** is used in adults to:

- relieve symptoms and heal duodenal ulcers.
- relieve symptoms and heal stomach ulcers.
- heal gastroesophageal reflux disease (GERD) and relieve its symptoms such as:
  - the burning feeling that rises from the chest to the throat (heartburn).
  - the flow of bitter/sour juice into the mouth (regurgitation).
- treat symptoms of non-erosive reflux disease (NERD) such as heartburn and regurgitation.
- maintain longer term healing of gastroesophageal reflux disease (GERD).
- treat rare conditions where excess acid is produced in the stomach (e.g., Zollinger-Ellison syndrome).
- treat ulcers caused by infection with the bacterium, *Helicobacter pylori* (*H. pylori*), and prevent these ulcers from coming back by:
  - taking **PARIET<sup>®</sup>** with antibiotics such as amoxicillin and clarithromycin, as directed by your doctor.
  - following the information on the antibiotics provided to you by the pharmacist.

**How does PARIET<sup>®</sup> work?**

**PARIET<sup>®</sup>** is a medicine called a proton pump inhibitor (PPI). **PARIET<sup>®</sup>** works by reducing the amount of acid made in your stomach.

**What are the ingredients in PARIET<sup>®</sup>?**

Medicinal ingredients: rabeprazole sodium

Non-medicinal ingredients: mannitol, hydroxypropyl cellulose, magnesium oxide, low-substituted hydroxypropyl cellulose, magnesium stearate, ethylcellulose, hydroxypropyl methylcellulose phthalate, diacetylated monoglycerides, talc, titanium dioxide, iron oxide (black, red, or yellow), and carnauba wax.

The 20 mg tablet also contains glycerine fatty acid ester.

**PARIET<sup>®</sup> comes in the following dosage forms:**

Tablets of 10 mg and 20 mg

**Do not use PARIET<sup>®</sup> if you are allergic to:**

- rabeprazole or other medications in this class.
- the “nonmedicinal” ingredients in **PARIET<sup>®</sup>** tablets (see **What are the ingredients in PARIET<sup>®</sup>?**).

- the antibiotics amoxicillin or clarithromycin if these are used with PARIET<sup>®</sup> to treat ulcers caused by *H. pylori*.

**Do not use PARIET<sup>®</sup> if you are:**

- taking rilpivirine

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take PARIET<sup>®</sup>. Talk about any health conditions or problems you may have, including:**

- all health problems you have now or have had in the past
- any liver problems
- if you experience symptoms including palpitations (rapid heartbeat), dizziness, seizures, twitching, spasms, muscle weakness, cramps and convulsions. These may be signs of low magnesium levels in your blood
- if you are taking methotrexate
- if you are pregnant or planning to become pregnant
- if you are breast-feeding or planning to breast-feed
- if you are due to have a specific blood test (Chromogranin A)

**Other warnings you should know about:**

Long-term use of PARIET<sup>®</sup> may prevent normal absorption of vitamin B<sub>12</sub> from the diet and could lead to vitamin B<sub>12</sub> deficiency. Talk to your doctor.

Using PARIET<sup>®</sup> for a long period of time (a year or longer) may increase the risk of fractures of the hip, wrist or spine. Talk to your doctor about your risk.

Using medicines like PARIET<sup>®</sup> for a long period of time may cause a growth in your stomach (polyp). This can lead to intestinal blockage or bleeding. Talk to your doctor if you experience nausea or stomach pain while you are taking PARIET<sup>®</sup>.

**Tell your healthcare professional about all the medicines you take, including any drugs (prescription and non-prescription), vitamins, minerals, natural supplements or alternative medicines.**

**The following may interact with PARIET<sup>®</sup>:**

- ketoconazole
- digoxin
- warfarin
- antiretroviral drugs such as atazanavir, nelfinavir, saquinavir
- methotrexate

**How to take PARIET<sup>®</sup>:**

- Take PARIET<sup>®</sup> exactly as prescribed by your doctor, usually for a specific number of weeks.
- Use the lowest dose and shortest time as determined by your doctor.
- Do not stop taking PARIET<sup>®</sup> even when you start to feel better. If you stop taking PARIET<sup>®</sup> too soon, your symptoms may return.

- Take PARIET<sup>®</sup> with or without meals.
- Swallow tablets whole with a liquid. Do not chew or crush the tablets.
- Talk to your doctor if you have any concerns.

#### Usual dose:

Condition	Adult Dose	How often	How long
reflux symptoms with esophagitis	20 mg	once daily	four weeks
reflux symptoms without esophagitis, such as heartburn and regurgitation	10 mg to a maximum of 20 mg	once daily	four weeks
duodenal ulcer	20 mg	once daily	Up to four weeks
stomach ulcer	20 mg	once daily	Up to six weeks
ulcer caused by <i>H. pylori</i> infection <sup>†</sup>	20 mg	twice daily in combination with antibiotic drugs (clarithromycin and amoxicillin) preferably with the morning and evening meals	one week

<sup>†</sup>If you are given PARIET<sup>®</sup> in combination with antibiotic drugs, it is important that you take all medications at the correct time of day and for the entire treatment period to ensure they will work properly. Studies have shown that patients who take their medications as prescribed have better ulcer healing rates and greater success in getting rid of their *H. pylori* infection.

#### Overdose:

If you think you have taken too much PARIET<sup>®</sup>, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

#### Missed Dose:

If you forget to take one dose of PARIET<sup>®</sup> medication, take a tablet as soon as you remember, unless it is almost time for your next dose. If it is, do not take the missed tablet at all. **Never double-up on a dose to make up for the one you have missed; just go back to your regular schedule.**

#### What are possible side effects from using PARIET<sup>®</sup>?

Like all medicines, PARIET<sup>®</sup> can cause side effects.

Headache and diarrhea are the most common side effects experienced with PARIET<sup>®</sup>. Less common side effects are rash, itchiness and dizziness. If any of these become troublesome, consult your doctor.

Treatment in combination with antibiotics: If you experience symptoms such as severe diarrhea (bloody or watery) with or without fever, abdominal pain, or tenderness, you may have

*Clostridium difficile* colitis (bowel inflammation). If this occurs, stop taking PARIET® and contact your healthcare professional immediately.

If you experience symptoms of low magnesium levels in your body, your doctor may stop PARIET®.

Tell your doctor right away if you have any of these symptoms:

- seizures.
- dizziness.
- abnormal or fast heart beat.
- jitteriness.
- jerking movements or shaking (tremors).
- muscle weakness.
- spasms of the hands and feet.
- cramps or muscle aches.
- spasm of the voice box.
- fracture (broken bone).
- New or worsening joint pain.
- Rash on your cheeks or arms that gets worse in the sun.
- Blood in stool.

Stopping your PPI therapy after taking it for a long time, may cause your symptoms to get worse and your stomach may increase acid production. Carefully follow your doctor's instructions when discontinuing your PPI therapy.

### **Reporting Side Effects**

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

#### **3 ways to report:**

- Online at MedEffect®  
([www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html](http://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html));
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
  - Fax to 1-866-678-6789 (toll-free), or
  - Mail to: Canada Vigilance Program  
Health Canada, Postal Locator 1908C  
Ottawa, ON  
K1A 0K9Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect® ([www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html](http://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html)).

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**Storage:**

Keep your tablets stored at room temperature (15° to 25°C) and protect from moisture. Keep out of the sight and reach of children.

**If you want more information about PARIET®:**

- Talk to your healthcare professional.
- For questions or concerns contact the manufacturer, Janssen Inc. ([www.janssen.com/canada](http://www.janssen.com/canada)).
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website [www.janssen.com/canada](http://www.janssen.com/canada), or by calling 1-800-567-3331 or 1-800-387-8781.

This leaflet was prepared by Janssen Inc. Toronto, Ontario, M3C 1L9.

Last Revised: November 2018